arXiv:math/0405106v1 [math.OA] 6 May 2004

AMALGAMATED R-TRANSFORM THEORY ON TOEPLITZ-PROBABILITY SPACES OVER TOEPLITZ MATRICIAL ALGEBRAS

ILWOO CHO

Abstract. In this paper, we will consider a noncommutative probability space, (T, E), over a Toeplitz matricial algebra $B = \mathcal{C}^N$, for $N \in \mathbb{N}$, induced by a (scalar-valued) noncommutative probability space, (A, φ), with a suitable B-functional, $E : T \to B$, defined by φ. On this framework, we will observe amalgamated R-transform calculus with respect to a B-functional, E. The technique and idea are came from those in [12] and Free Probability of type B studied in [31].

Voiculescu developed Free Probability Theory. Here, the classical concept of Independence in Probability theory is replaced by a noncommutative analogue called Freeness (See [9]). There are two approaches to study Free Probability Theory. One of them is the original analytic approach of Voiculescu and the other one is the combinatorial approach of Speicher and Nica (See [1], [2] and [3]).

To observe the free additive convolution and free multiplicative convolution of two distributions induced by free random variables in a noncommutative probability space (over $B = \mathbb{C}$), Voiculescu defined the R-transform and the S-transform, respectively. These show that to study distributions is to study certain (B-)formal series for arbitrary noncommutative indeterminants.

Speicher defined the free cumulants which are the main objects in the combinatorial approach of Free Probability Theory. And he developed free probability theory by using Combinatorics and Lattice theory on collections of noncrossing partitions (See [3]). Also, Speicher considered the operator-valued free probability theory, which is also defined and observed analytically by Voiculescu, when $\mathbb{C}$ is replaced to an arbitrary algebra B (See [1]). Nica defined R-transforms of several random variables (See [2]). He defined these R-transforms as multivariable formal series in noncommutative several indeterminants. To observe the R-transform, the Möbius Inversion under the embedding of lattices plays a key role (See [1],[3],[5],[12],[13] and [17]).

In [12], [19] and [20], we observed the amalgamated R-transform calculus. Actually, amalgamated R-transforms are defined originally by Voiculescu (See [10]) and are characterized combinatorially by Speicher (See [1]). In [12], we defined amalgamated R-transforms slightly differently from those defined in [1] and [10]. We defined them as B-formal series and tried to characterize, like in [2] and [3]. The main tool which is considered, for studying amalgamated R-transforms is amalgamated boxed convolution. After defining boxed convolution over an arbitrary algebra B, we could get that

$$R_{x_1,...,x_s} \boxed{*} R_{y_1,...,y_s}^{symm(1_B)} = R_{x_1y_1,...,x_sy_s}, \text{ for any } s \in \mathbb{N},$$

where x_j and y_j are free B-valued random variables.

In this paper, we will consider a noncommutative probability space, (T, E), over a Toeplitz matricial algebra $B = \mathcal{C}^N$, for $N \in \mathbb{N}$, induced by a (scalar-valued) noncommutative probability space, (A, φ), with a suitable B-functional, $E : T \to B$, defined by φ. On this framework, we will observe amalgamated R-transform calculus with respect to a B-functional, E. The technique and idea are came from those in [12] and Free Probability of type B studied in [31]. By the construction of this amalgamated noncommutative probability space, we can easily recognize that to compute the partition-dependent moments of random variables, we do not need to concern about the insertion property. We will concentrate on observing the moment series and R-transforms of random variables in this structure and after considering the (combinatorial)amalgamated freeness, we construct the R-transform calculus.

1. Preliminaries

1.1. **Free Probability Theory.**

In this section, we will summarize and introduced the basic results from [1] and [12]. Throughout this section, let B be a unital algebra. The algebraic pair (A, φ) is said to be a noncommutative probability space over B (shortly, NCPSpace over B) if A is an algebra over B (i.e $1_B = 1_A \in B \subset A$) and $\varphi : A \to B$ is a B-functional (or a conditional expectation) ; φ satisfies

$$\varphi(b) = b, \text{ for all } b \in B$$

and

$$\varphi(bxb') = b\varphi(x)b', \text{ for all } b, b' \in B \text{ and } x \in A.$$

Let (A, φ) be a NCPSpace over B. Then, for the given B-functional, we can determine a moment multiplicative function $\widehat{\varphi} = (\varphi^{(n)})_{n=1}^{\infty} \in I(A, B)$, where

$$\varphi^{(n)}(a_1 \otimes ... \otimes a_n) = \varphi(a_1....a_n),$$

for all $a_1 \otimes ... \otimes a_n \in A^{\otimes_B n}$, $\forall n \in \mathbb{N}$.

We will denote noncrossing partitions over $\{1, ..., n\}$ ($n \in \mathbb{N}$) by $NC(n)$. Define an ordering on $NC(n)$;

$\theta = \{V_1, ..., V_k\} \leq \pi = \{W_1, ..., W_l\} \overset{def}{\Leftrightarrow}$ For each block $V_j \in \theta$, there exists only one block $W_p \in \pi$ such that $V_j \subset W_p$, for $j = 1, ..., k$ and $p = 1, ..., l$.

Then $(NC(n), \leq)$ is a complete lattice with its minimal element $0_n = \{(1), ..., (n)\}$ and its maximal element $1_n = \{(1, ..., n)\}$. We define the incidence algebra I_2 by a set of all complex-valued functions η on $\cup_{n=1}^{\infty} (NC(n) \times NC(n))$ satisfying $\eta(\theta, \pi) = 0$, whenever $\theta \nleq \pi$. Then, under the convolution

$$* : I_2 \times I_2 \rightarrow \mathbb{C}$$

defined by

$$\eta_1 * \eta_2(\theta, \pi) = \sum_{\theta \leq \sigma \leq \pi} \eta_1(\theta, \sigma) \cdot \eta_2(\sigma, \pi),$$

I_2 is indeed an algebra of complex-valued functions. Denote zeta, Möbius and delta functions in the incidence algebra I_2 by ζ, μ and δ, respectively. i.e

$$\zeta(\theta, \pi) = \begin{cases} 1 & \theta \leq \pi \\ 0 & otherwise, \end{cases}$$

$$\delta(\theta, \pi) = \begin{cases} 1 & \theta = \pi \\ 0 & otherwise, \end{cases}$$

and μ is the $(*)$-inverse of ζ. Notice that δ is the $(*)$-identity of I_2. By using the same notation $(*)$, we can define a convolution between $I(A, B)$ and I_2 by

$$\widehat{f} * \eta \, (a_1, ..., a_n \, ; \, \pi) = \sum_{\pi \in NC(n)} \widehat{f}(\pi)(a_1 \otimes ... \otimes a_n)\eta(\pi, 1_n),$$

where $\widehat{f} \in I(A, B)$, $\eta \in I_1$, $\pi \in NC(n)$ and $a_j \in A$ ($j = 1, ..., n$), for all $n \in \mathbb{N}$. Notice that $\widehat{f} * \eta \in I(A, B)$, too. Let $\widehat{\varphi}$ be a moment multiplicative function in $I(A, B)$ which we determined before. Then we can naturally define a cumulant multiplicative function $\widehat{c} = (c^{(n)})_{n=1}^{\infty} \in I(A, B)$ by

$$\widehat{c} = \widehat{\varphi} * \mu \quad \text{or} \quad \widehat{\varphi} = \widehat{c} * \zeta.$$

This says that if we have a moment multiplicative function, then we always get a cumulant multiplicative function and vice versa, by $(*)$. This relation is so-called "Möbius Inversion". More precisely, we have

$$\begin{aligned}\varphi(a_1 ... a_n) &= \varphi^{(n)}(a_1 \otimes ... \otimes a_n) \\ &= \sum_{\pi \in NC(n)} \widehat{c}(\pi)(a_1 \otimes ... \otimes a_n)\zeta(\pi, 1_n) \\ &= \sum_{\pi \in NC(n)} \widehat{c}(\pi)(a_1 \otimes ... \otimes a_n),\end{aligned}$$

for all $a_j \in A$ and $n \in \mathbb{N}$. Or equivalently,

$$c^{(n)}(a_1 \otimes ... \otimes a_n) = \sum_{\pi \in NC(n)} \widehat{\varphi}(\pi)(a_1 \otimes ... \otimes a_n)\mu(\pi, 1_n).$$

Now, let (A_i, φ_i) be NCPSpaces over B, for all $i \in I$. Then we can define a amalgamated free product of A_i 's and amalgamated free product of φ_i's by

$$A \equiv *_B A_i \quad \text{and} \quad \varphi \equiv *_i \varphi_i,$$

respectively. Then, by Voiculescu, (A, φ) is again a NCPSpace over B and, as a vector space, A can be represented by

$$A = B \oplus \left(\oplus_{n=1}^{\infty} \left(\underset{i_1 \neq ... \neq i_n}{\oplus} (A_{i_1} \ominus B) \otimes ... \otimes (A_{i_n} \ominus B) \right) \right),$$

where $A_{i_j} \ominus B = \ker \varphi_{i_j}$. We will use Speicher's combinatorial definition of amalgamated free product of B-functionals ;

Definition 1.1. *Let (A_i, φ_i) be NCPSpaces over B, for all $i \in I$. Then $\varphi = *_i \varphi_i$ is the amalgamated free product of B-functionals φ_i's on $A = *_B A_i$ if the cumulant multiplicative function $\widehat{c} = \widehat{\varphi} * \mu \in I(A, B)$ has its restriction to $\underset{i \in I}{\cup} A_i$, $\underset{i \in I}{\oplus} \widehat{c}_i$, where $\widehat{c}_i$ is the cumulant multiplicative function induced by φ_i, for all $i \in I$. i.e*

$$c^{(n)}(a_1 \otimes ... \otimes a_n) = \begin{cases} c_i^{(n)}(a_1 \otimes ... \otimes a_n) & \text{if } \forall a_j \in A_i \\ 0_B & \text{otherwise.} \end{cases}$$

Now, we will observe the freeness over B ;

Definition 1.2. *Let (A, φ)be a NCPSpace over B.*

(1) Subalgebras containing B, $A_i \subset A$ $(i \in I)$ are free (over B) if we let $\varphi_i = \varphi \mid_{A_i}$, for all $i \in I$, then $*_i \varphi_i$ has its cumulant multiplicative function $\widehat{c}$ such that its restriction to $\underset{i \in I}{\cup} A_i$ is $\underset{i \in I}{\oplus} \widehat{c}_i$, where $\widehat{c}_i$ is the cumulant multiplicative function induced by each φ_i, for all $i \in I$.

(2) Sebsets X_i $(i \in I)$ are free (over B) if subalgebras A_i's generated by B and X_i's are free in the sense of (1). i.e If we let $A_i = A\lg(X_i, B)$, for all $i \in I$, then A_i's are free over B.

In [1], Speicher showed that the above combinatorial freeness with amalgamation can be used alternatively with respect to Voiculescu's original freeness with amalgamation.

Let (A, φ) be a NCPSpace over B and let $x_1, ..., x_s$ be B-valued random variables $(s \in \mathbb{N})$. Define $(i_1, ..., i_n)$-th moment of $x_1, ..., x_s$ by

$$\varphi(x_{i_1} b_{i_2} x_{i_2} ... b_{i_n} x_{i_n}),$$

for arbitrary $b_{i_2}, ..., b_{i_n} \in B$, where $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$. Similarly, define a symmetric $(i_1, ..., i_n)$-th moment by the fixed $b_0 \in B$ by

$$\varphi(x_{i_1} b_0 x_{i_2} ... b_0 x_{i_n}).$$

If $b_0 = 1_B$, then we call this symmetric moments, trivial moments.

Cumulants defined below are main tool of combinatorial free probability theory ; in [12], we defined the $(i_1, ..., i_n)$-th cumulant of $x_1, ..., x_s$ by

$$k_n(x_{i_1}, ..., x_{i_n}) = c^{(n)}(x_{i_1} \otimes b_{i_2} x_{i_2} \otimes ... \otimes b_{i_n} x_{i_n}),$$

for $b_{i_2}, ..., b_{i_n} \in B$, arbitrary, and $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$, where $\widehat{c} = (c^{(n)})_{n=1}^{\infty}$ is the cumulant multiplicative function induced by φ. Notice that, by Möbius inversion, we can always take such B-value whenever we have $(i_1, ..., i_n)$-th moment of $x_1, ..., x_s$. And, vice versa, if we have cumulants, then we can always take moments. Hence we can define a symmetric $(i_1, ..., i_n)$-th cumulant by $b_0 \in B$ of $x_1, ..., x_s$ by

$$k_n^{symm(b_0)}(x_{i_1}, ..., x_{i_n}) = c^{(n)}(x_{i_1} \otimes b_0 x_{i_2} \otimes ... \otimes b_0 x_{i_n}).$$

If $b_0 = 1_B$, then it is said to be trivial cumulants of $x_1, ..., x_s$.

By Speicher, it is shown that subalgebras A_i $(i \in I)$ are free over B if and only if all mixed cumulants vanish.

Proposition 1.1. *(See [1] and [12]) Let (A, φ) be a NCPSpace over B and let $x_1, ..., x_s \in (A, \varphi)$ be B-valued random variables $(s \in \mathbb{N})$. Then $x_1, ..., x_s$ are free if and only if all their mixed cumulants vanish.* $\square$

1.2. **Amalgamated R-transform Theory.**

In this section, we will define an R-transform of several B-valued random variables. Note that to study R-transforms is to study operator-valued distributions. R-transforms with single variable is defined by Voiculescu (over B, in particular,

$B = \mathbb{C}$. See [9] and [10]). Over $\mathbb{C}$, Nica defined multi-variable R-transforms in [2]. In [12], we extended his concepts, over B. R-transforms of B-valued random variables can be defined as B-formal series with its $(i_1, ..., i_n)$-th coefficients, $(i_1, ..., i_n)$-th cumulants of B-valued random variables, where $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$.

Definition 1.3. *Let (A, φ) be a NCPSpace over B and let $x_1, ..., x_s \in (A, \varphi)$ be B-valued random variables ($s \in \mathbb{N}$). Let $z_1, ..., z_s$ be noncommutative indeterminants. Define a moment series of $x_1, ..., x_s$, as a B-formal series, by*

$$M_{x_1,...,x_s}(z_1, ..., z_s) = \sum_{n=1}^{\infty} \sum_{i_1,..,i_n \in \{1,...,s\}} \varphi(x_{i_1} b_{i_2} x_{i_2} ... b_{i_n} x_{i_n})\, z_{i_1} ... z_{i_n},$$

where $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary for all $(i_2, ..., i_n) \in \{1, ..., s\}^{n-1}$, $\forall n \in \mathbb{N}$.

Define an R-transform of $x_1, ..., x_s$, as a B-formal series, by

$$R_{x_1,...,x_s}(z_1, ..., z_s) = \sum_{n=1}^{\infty} \sum_{i_1,...,i_n \in \{1,...,s\}} k_n(x_{i_1}, ..., x_{i_n})\, z_{i_1} ... z_{i_n},$$

with

$$k_n(x_{i_1}, ..., x_{i_n}) = c^{(n)}(x_{i_1} \otimes b_{i_2} x_{i_2} \otimes ... \otimes b_{i_n} x_{i_n}),$$

where $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary for all $(i_2, ..., i_n) \in \{1, ..., s\}^{n-1}$, $\forall n \in \mathbb{N}$. Here, $\widehat{c} = (c^{(n)})_{n=1}^{\infty}$ is a cumulant multiplicative function induced by φ in $I(A, B)$.

Denote a set of all B-formal series with s-noncommutative indeterminants ($s \in \mathbb{N}$), by Θ_B^s. i.e if $g \in \Theta_B^s$, then

$$g(z_1, ..., z_s) = \sum_{n=1}^{\infty} \sum_{i_1,...,i_n \in \{1,...,s\}} b_{i_1,...,i_n}\, z_{i_1} ... z_{i_n},$$

where $b_{i_1,...,i_n} \in B$, for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$. Trivially, by definition, $M_{x_1,...,x_s}$, $R_{x_1,...,x_s} \in \Theta_B^s$. By $\mathcal{R}_B^s$, we denote a set of all R-transforms of s-B-valued random variables. Recall that, set-theoratically,

$$\Theta_B^s = \mathcal{R}_B^s, \text{ sor all } s \in \mathbb{N}.$$

By the previous section, we can also define symmetric moment series and symmetric R-transform by $b_0 \in B$, by

$$M_{x_1,...,x_s}^{symm(b_0)}(z_1, ..., z_s) = \sum_{n=1}^{\infty} \sum_{i_1,...,i_n \in \{1,...,s\}} \varphi(x_{i_1} b_0 x_{i_2} ... b_0 x_{i_n})\, z_{i_1} ... z_{i_n}$$

and

$$R_{x_1,...,x_s}^{symm(b_0)}(z_1, ..., z_s) = \sum_{n=1}^{\infty} \sum_{i_1,..,i_n \in \{1,...,s\}} k_n^{symm(b_0)}(x_{i_1}, ..., x_{i_n})\, z_{i_1} ... z_{i_n},$$

with

$$k_n^{symm(b_0)}(x_{i_1}, ..., x_{i_n}) = c^{(n)}(x_{i_1} \otimes b_0 x_{i_2} \otimes ... \otimes b_0 x_{i_n}),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$.

If $b_0 = 1_B$, then we have trivial moment series and trivial R-transform of $x_1, ..., x_s$ denoted by $M^t_{x_1,...,x_s}$ and $R^t_{x_1,...,x_s}$, respectively.

The followings are known in [1] and [12] ;

Proposition 1.2. *Let (A, φ) be a NCPSpace over B and let $x_1, ..., x_s, y_1, ..., y_p \in (A, \varphi)$ be B-valued random variables, where $s, p \in \mathbb{N}$. Suppose that $\{x_1, ..., x_s\}$ and $\{y_1, ..., y_p\}$ are free in (A, φ). Then*

(1) $R_{x_1,...,x_s,y_1,...,y_p}(z_1, ..., z_{s+p}) = R_{x_1,...,x_s}(z_1, ..., z_s) + R_{y_1,...,y_p}(z_1, ..., z_p)$.

(2) If $s = p$, then $R_{x_1+y_1,...,x_s+y_s}(z_1, ..., z_s) = \left(R_{x_1,...,x_s} + R_{y_1,...,y_s}\right)(z_1, ..., z_s)$.
$\square$

The above proposition is proved by the characterization of freeness with respect to cumulants. i.e $\{x_1, ..., x_s\}$ and $\{y_1, ..., y_p\}$ are free in (A, φ) if and only if their mixed cumulants vanish. Thus we have

$$
\begin{aligned}
&k_n(p_{i_1}, ..., p_{i_n}) = c^{(n)}(p_{i_1} \otimes b_{i_2}p_{i_2} \otimes ... \otimes b_{i_n}p_{i_n}) \\
&\quad = \left(\widehat{c_x} \oplus \widehat{c_y}\right)^{(n)} \left(p_{i_1} \otimes b_{i_2}p_{i_2} \otimes ... \otimes b_{i_n}p_{i_n}\right) \\
&\quad = \left\{ \begin{array}{ll} k_n(x_{i_1}, ..., x_{i_n}) & \quad or \\ k_n(y_{i_1}, ..., y_{i_n}) & \end{array} \right.
\end{aligned}
$$

and if $s = p$, then

$$
\begin{aligned}
&k_n(x_{i_1}+y_{i_1}, ..., x_{i_n}+y_{i_n}) = c^{(n)}\left((x_{i_1} + y_{i_1}) \otimes b_{i_2}(x_{i_2} + y_{i_2}) \otimes ... \otimes b_{i_n}(x_{i_n} + y_{i_n})\right) \\
&\quad = c^{(n)}(x_{i_1} \otimes b_{i_2}x_{i_2} \otimes ... \otimes b_{i_n}x_{i_n}) + c^{(n)}(y_{i_1} \otimes b_{i_2}y_{i_2} \otimes ... \otimes b_{i_n}y_{i_n}) + [Mixed]
\end{aligned}
$$

where $[Mixed]$ is the sum of mixed cumulants, by the bimodule map property of $c^{(n)}$

$$= k_n(x_{i_1}, ..., x_{i_n}) + k_n(y_{i_1}, ..., y_{i_n}) + 0_B.$$

Now, we will define B-valued boxed convolution $\boxed{*}$, as a binary operation on Θ^s_B ; note that if $f, g \in \Theta^s_B$, then we can always choose free $\{x_1, ..., x_s\}$ and $\{y_1, ..., y_s\}$ in (some) NCPSpace over B, (A, φ), such that

$$f = R_{x_1,...,x_s} \quad \text{and} \quad g = R_{y_1,...,y_s}.$$

Definition 1.4. *(1) Let $s \in \mathbb{N}$. Let $(f, g) \in \Theta^s_B \times \Theta^s_B$. Define $\boxed{*} : \Theta^s_B \times \Theta^s_B \to \Theta^s_B$ by*

$$(f, g) = (R_{x_1,...,x_s},\, R_{y_1,...,y_s}) \longmapsto R_{x_1,...,x_s} \,\boxed{*}\, R_{y_1,...,y_s}.$$

Here, $\{x_1, ..., x_s\}$ and $\{y_1, ..., y_s\}$ are free in (A, φ). Suppose that

$$coef_{i_1,..,i_n}\left(R_{x_1,...,x_s}\right) = c^{(n)}(x_{i_1} \otimes b_{i_2}x_{i_2} \otimes ... \otimes b_{i_n}x_{i_n})$$

and

$$coef_{i_1,...,i_n}(R_{y_1,...,y_s}) = c^{(n)}(y_{i_1} \otimes b'_{i_2}y_{i_2} \otimes ... \otimes b'_{i_n}y_{i_n}),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$, *where* $b_{i_j}, b'_{i_n} \in B$ *arbitrary. Then*

$$coef_{i_1,...,i_n}\left(R_{x_1,...,x_s} \boxed{*} R_{y_1,...,y_s}\right)$$

$$= \sum_{\pi \in NC(n)} (\widehat{c_x} \oplus \widehat{c_y})\,(\pi \cup Kr(\pi))(x_{i_1} \otimes y_{i_1} \otimes b_{i_2}x_{i_2} \otimes b'_{i_2}y_{i_2} \otimes ... \otimes b_{i_n}x_{i_n} \otimes b'_{i_n}y_{i_n})$$

$$\overset{denote}{=} \sum_{\pi \in NC(n)} \left(k_\pi \oplus k_{Kr(\pi)}\right)(x_{i_1}, y_{i_1}, ..., x_{i_n}y_{i_n}),$$

where $\widehat{c_x} \oplus \widehat{c_y} = \widehat{c}\mid_{A_x *_B A_y}$, $A_x = A\lg(\{x_i\}_{i=1}^s, B)$ *and* $A_y = A\lg(\{y_i\}_{i=1}^s, B)$ *and where* $\pi \cup Kr(\pi)$ *is an alternating union of partitions in* $NC(2n)$

Proposition 1.3. *(See [12])Let* (A, φ) *be a NCPSpace over* B *and let* $x_1, ..., x_s, y_1, ..., y_s \in (A, \varphi)$ *be* B*-valued random variables* ($s \in \mathbb{N}$). *If* $\{x_1, ..., x_s\}$ *and* $\{y_1, ..., y_s\}$ *are free in* (A, φ), *then we have*

$$k_n(x_{i_1}y_{i_1}, ..., x_{i_n}y_{i_n})$$

$$= \sum_{\pi \in NC(n)} (\widehat{c_x} \oplus \widehat{c_y})\,(\pi \cup Kr(\pi))(x_{i_1} \otimes y_{i_1} \otimes b_{i_2}x_{i_2} \otimes y_{i_2} \otimes ... \otimes b_{i_n}x_{i_n} \otimes y_{i_n})$$

$$\overset{denote}{=} \sum_{\pi \in NC(n)} \left(k_\pi \oplus k_{Kr(\pi)}^{symm(1_B)}\right)(x_{i_1}, y_{i_1}, ..., x_{i_n}, y_{i_n}),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$, $b_{i_2}, ..., b_{i_n} \in B$, *arbitrary, where* $\widehat{c_x} \oplus \widehat{c_y} = \widehat{c}\mid_{A_x *_B A_y}$, $A_x = A\lg(\{x_i\}_{i=1}^s, B)$ *and* $A_y = A\lg(\{y_i\}_{i=1}^s, B)$. $\square$

This shows that ;

Corollary 1.4. *(See [12]) Under the same condition with the previous proposition,*

$$R_{x_1,...,x_s} \boxed{*} R^t_{y_1,...,y_s} = R_{x_1y_1,...,x_sy_s}.$$

$\square$

Notice that, in general, unless $b'_{i_2} = ... = b'_{i_n} = 1_B$ in B,

$$R_{x_1,...,x_s} \boxed{*} R_{y_1,...,y_s} \neq R_{x_1y_1,...,x_sy_s}.$$

However, as we can see above,

$$R_{x_1,...,x_s} \boxed{*} R^t_{y_1,...,y_s} = R_{x_1y_1,...,x_sy_s}$$

and

$$R^t_{x_1,...,x_s} \boxed{*} R^t_{y_1,...,y_s} = R^t_{x_1y_1,...,x_sy_s},$$

where $\{x_1, ..., x_s\}$ and $\{y_1, ..., y_s\}$ are free over B. Over $B = \mathbb{C}$, the last equation is proved by Nica and Speicher in [2] and [3]. Actually, their R-transforms (over $\mathbb{C}$) is our trivial R-transforms (over $\mathbb{C}$). Is it possible to find a trivial R-transform $R^t_{x'_1,...,x'_s}$, for the given R-transform $R_{x_1,...,x_s}$, where $x_1, ..., x_s \in (A, \varphi)$ are B-valued random variables and $x'_1, ..., x'_s$ are B-valued random variables (which are not necessarily contained in (A, φ)) ? If possible, to study amalgamated R-transforms, we can only deal with trivial R-transforms. Let's denote a set of all tritial R-transforms by $S^{(1_B)}\,\mathcal{R}^s_B$. If we have a positive answer of above question, then $\mathcal{R}^s_B = S^{(1_B)}\,\mathcal{R}^s_B$. In that case, we can deal with B-valued boxed convolution more freely. In the following chapter, we will consider this possibility.

1.3. **Scalar-Valued Case (in the sense of Nica and Speicher).**

Suppose that $B = \mathbb{C}$. Then a NCPSpace over B, (A, φ), All information of B-valued (scalar-valued) distributions are contained in corresponding trivial R-transforms (See [32] and [33]). So, WLOG, we can define an R-transform of random variable(s) as a trivial R-transform of the random variable(s). This is just an (scalar-valued) R-transform in the sense of Nica (See [2]). In [32] and [33], we showed that if B is an arbitrary algebra such that $B = C_A(B)$, then amalgamated freeness can be characterized by

$$\text{Amalgamated Freeness} \overset{def}{\Leftrightarrow} \text{All mixed cumulants vanish}$$
$$\Leftrightarrow \text{All mixed } \textbf{trivial} \text{ cumulants vanish.}$$

Since $C_A(\mathbb{C}) = \mathbb{C}$, for any algebra A, if $B = \mathbb{C}$, then the freeness is characterized by trivial cumulants.

Definition 1.5. *Let (A, φ) be a noncommutative probability space over $\mathbb{C}$ (shortly NCPSpace) with a linear functional $\varphi : A \to \mathbb{C}$ and let $x_1, ..., x_s \in (A, \varphi)$ be (scalar-valued) random variables ($s \in \mathbb{N}$). Define a moment series and R-transform of $x_1, ..., x_s$ by $M^t_{x_1,...,x_s}$ and $R^t_{x_1,...,x_s}$, respectively, where $M^t_{x_1,...,x_s}$ and $R^t_{x_1,...,x_s}$ are defined in Section 1.2. By $M_{x_1,...,x_s}$ and $R_{x_1,...,x_s}$, we will denote the moment series and the R-transform, respectively. Also, if we mention cumulants, then they are trivial cumulants, just denoted by $k_n(\cdot)$, for $n \in \mathbb{N}$.*

Remark 1.1. *By definition, all facts, which we considered in the previous sections, holds true, for new* $M_{x_1,...,x_s}$ *and* $R_{x_1,...,x_s}$*. Moreover, if* $\{x_1, ..., x_s\}$ *and* $\{y_1, ..., y_s\}$ *are free subsets of random variables, then*

$$R_{x_1y_1,...,x_sy_s} = R_{x_1,...,x_s} \boxtimes_{\mathbb{C}} R_{y_1,...,y_s},$$

where $\boxtimes_{\mathbb{C}}$ *is the* $B = \mathbb{C}$*-valued boxed convolution. This is the boxed convolution* $\boxtimes$ *in the sense of Nica (See [2]). Notice that all random variables are central over* $\mathbb{C}$*. So, we have that*

$$\begin{aligned} &coef_{i_1,...,i_n}\left(R_{x_1,...,x_s} \boxtimes R_{y_1,...,y_s}\right) \\ &\qquad = \sum_{\pi \in NC(n)} \left(k_\pi(x_{i_1}, ..., x_{i_n})\right)\left(k_{Kr(\pi)}(y_{i_1}, ..., y_{i_n})\right). \end{aligned}$$

Again, we can see that this is just a definition of boxed convolution in the sense of Nica and Speicher (See [2] and [3]).

Form now, by (A, φ), we will denote a NCPSpace.

2. Toeplitz Matricial Probability Theory

2.1. **Toeplitz Matricial Probability Spaces over Toeplitz Matricial Algebras.**

In this section, we will consider the Toeplitz Matricial probability space over a Toeplitz Matricial Algebra $\mathcal{C}^n \cong \mathbb{C}^n$, for $n \in \mathbb{N}$. We define a Toeplitz Matricial algebra $\mathcal{C}^n$ by an algebra $\mathbb{C}^n$ with following addition and multiplication ;

$$(\alpha_1, ..., \alpha_n) + (\alpha_1', ..., \alpha_n') = (\alpha_1 + \alpha_1', ..., \alpha_n + \alpha_n')$$

and

$$(\alpha_1, ..., \alpha_n) \cdot (\alpha_1', ..., \alpha_n') = (\alpha_1\alpha_1', \textstyle\sum_{k=1}^{2} \alpha_k \alpha'_{(2+1)-k}, ..., \sum_{k=1}^{n} \alpha_k \alpha'_{(n+1)-k}),$$

respectively. Then $\mathcal{C}^n = (\mathbb{C}^n, +, \cdot)$ is an algebra. And this algebra is called an n-th Toeplitz Matricial algebra. i.e isometrically, this algebra $\mathcal{C}^n$ can be understood as an algebra generated by all Toeplitz matrices in $M_n(\mathbb{C})$ which have the form of

$$(\alpha_1, ..., \alpha_n) = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_n \\ & \alpha_1 & \alpha_2 & \ddots & \vdots \\ & & \alpha_1 & \ddots & \alpha_3 \\ & & & \ddots & \alpha_2 \\ O & & & & \alpha_1 \end{pmatrix} \in M_n(\mathbb{C}),$$

for all $(\alpha_1, ..., \alpha_n) \in \mathcal{C}^n$, $n \in \mathbb{N}$.

For exmaple, if $n = 4$, then we have that

$$(\alpha_1, \alpha_2, \alpha_3, \alpha_4) + (\alpha_1', \alpha_2', \alpha_3', \alpha_4')$$

$$\cong \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ 0 & \alpha_1 & \alpha_2 & \alpha_3 \\ 0 & 0 & \alpha_1 & \alpha_2 \\ 0 & 0 & 0 & \alpha_1 \end{pmatrix} + \begin{pmatrix} \alpha_1' & \alpha_2' & \alpha_3' & \alpha_4' \\ 0 & \alpha_1' & \alpha_2' & \alpha_3' \\ 0 & 0 & \alpha_1' & \alpha_2' \\ 0 & 0 & 0 & \alpha_1' \end{pmatrix}$$

$$= \begin{pmatrix} \alpha_1 + \alpha_1' & \alpha_2 + \alpha_2' & \alpha_3 + \alpha_3' & \alpha_4 + \alpha_4' \\ 0 & \alpha_1 + \alpha_1' & \alpha_2 + \alpha_2' & \alpha_3 + \alpha_3' \\ 0 & 0 & \alpha_1 + \alpha_1' & \alpha_2 + \alpha_2' \\ 0 & 0 & 0 & \alpha_1 + \alpha_1' \end{pmatrix}$$

$$\cong (\alpha_1 + \alpha_1',\ \alpha_2 + \alpha_2',\ \alpha_3 + \alpha_3',\ \alpha_4 + \alpha_4')$$

and

$$(\alpha_1, \alpha_2, \alpha_3, \alpha_4) \cdot (\alpha_1', \alpha_2', \alpha_3', \alpha_4')$$

$$\cong \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ 0 & \alpha_1 & \alpha_2 & \alpha_3 \\ 0 & 0 & \alpha_1 & \alpha_2 \\ 0 & 0 & 0 & \alpha_1 \end{pmatrix} \begin{pmatrix} \alpha_1' & \alpha_2' & \alpha_3' & \alpha_4' \\ 0 & \alpha_1' & \alpha_2' & \alpha_3' \\ 0 & 0 & \alpha_1' & \alpha_2' \\ 0 & 0 & 0 & \alpha_1' \end{pmatrix}$$

$$= \begin{pmatrix} \alpha_1\alpha_1' & \alpha_1\alpha_2' + \alpha_2\alpha_1' & \alpha_1\alpha_3' + \alpha_2\alpha_2' + \alpha_3\alpha_1' & \alpha_1\alpha_4' + \alpha_2\alpha_3' + \alpha_3\alpha_2' + \alpha_4\alpha_1' \\ 0 & \alpha_1\alpha_1' & \alpha_1\alpha_2' + \alpha_2\alpha_1' & \alpha_1\alpha_3' + \alpha_2\alpha_2' + \alpha_3\alpha_1' \\ 0 & 0 & \alpha_1\alpha_1' & \alpha_1\alpha_2' + \alpha_2\alpha_1' \\ 0 & 0 & 0 & \alpha_1\alpha_1' \end{pmatrix}$$

$$\cong (\alpha_1\alpha_1',\ \alpha_1\alpha_2' + \alpha_2\alpha_1',\ \alpha_1\alpha_3' + \alpha_2\alpha_2' + \alpha_3\alpha_1',\ \alpha_1\alpha_4' + \alpha_2\alpha_3' + \alpha_3\alpha_2' + \alpha_4\alpha_1'),$$

for all $(\alpha_1, ..., \alpha_4)$, $(\alpha_1', ..., \alpha_4') \in \mathcal{C}^4$. Notice that $(1, 0, ..., 0) \in \mathcal{C}^n$ is the multiplication-identity on $\mathcal{C}^n$.

We will observe the R-transform calculus in a noncommutative probability space over a Toeplitz matricial algebra $\mathcal{C}^n$.

$\boxed{\text{Notation}}$ From now, for any $n \in \mathbb{N}$, by little abuse of notation, we will denote $\mathcal{C}^n$ by B, if there is no confusion. $\square$

Let (A, φ) be a NCPSpace in the sense of Section 1.3, where A is an algebra and $\varphi : A \to \mathbb{C}$ is a linear functional. Then we can define a Toeplitz matricial algebra T_n over a Toeplitz matricial algebra $B = \mathcal{C}^n$, induced by (A, φ) ;

Definition 2.1. *Fix $n \in \mathbb{N}$. Let $B = \mathcal{C}^n$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace in the sense of Section 1.3, with a linear functional $\varphi : A \to \mathbb{C}$. Define a Toeplitz Matricial Probability Space over B, (T_n, E_n), induced by (A, φ), by*

(i) T_n is an algebra over B (i.e $1_B = (1, 0, ..., 0) \in B \subset T_n$) such that

$$T_n = A\lg\left(\{(a_1, ..., a_n) : \forall\, a_j \in (A, \varphi)\}\right)$$

with

$$(a_1, ..., a_n) + (a_1', ..., a_n') = (a_1 + a_1', ..., a_n + a_n')$$

and

$$(a_1, ..., a_n) \cdot (a_1', ..., a_n') = (a_1 a_1',\, \sum_{k=1}^{2} a_k a'_{(2+1)-k}, ..., \sum_{k=1}^{n} a_k a'_{(n+1)-k}),$$

for all $(a_1, ..., a_n)$, $(a_1', ..., a_n') \in T_n$.

(ii) $E : T_n \to B$ is a B-functional

$$E\left((a_1, ..., a_n)\right) = \left(\varphi(a_1), ..., \varphi(a_n)\right).$$

Without confusion, we will denote (T_n, E_n) by (T, E).

Lemma 2.1. *Let $n \in \mathbb{N}$. A Toeplitz matricial probability space, (T, E), over $B = \mathcal{C}^n$, is, indeed, a NCPSPace over B.*

Proof. Since T is an algebra over B, by definition, it suffices to show that a map $E : T \to B$ is a B-functional.

(i) Let $(\alpha_1, ..., \alpha_n) \in \mathcal{C}^n = B$. Then

$$E\left((\alpha_1, ..., \alpha_n)\right) = (\varphi(\alpha_1), ..., \varphi(\alpha_n)) = (\alpha_1, ..., \alpha_n),$$

since φ is a linear functional.

(ii) Let $(\alpha_1, ..., \alpha_n), (\alpha_1', ..., \alpha_n') \in B$ and $(a_1, ..., a_n) \in T$. Then

$$
\begin{aligned}
&E\left((\alpha_1,...,\alpha_n)(a_1,...,a_n)(\alpha'_1,...,\alpha'_n)\right)\\
&\equiv E\left(\begin{pmatrix} \alpha_1 & \alpha_2 & \cdots & \alpha_n \\ & \alpha_1 & \ddots & \vdots \\ & & \ddots & \alpha_2 \\ O & & & \alpha_1 \end{pmatrix}\begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ & a_1 & \ddots & \vdots \\ & & \ddots & a_2 \\ O & & & a_1 \end{pmatrix}\begin{pmatrix} \alpha'_1 & \alpha'_2 & \cdots & \alpha'_n \\ & \alpha'_1 & \ddots & \vdots \\ & & \ddots & \alpha'_2 \\ & & & \alpha'_1 \end{pmatrix}\right)\\
&= \begin{pmatrix} \alpha_1 & \alpha_2 & \cdots & \alpha_n \\ & \alpha_1 & \ddots & \vdots \\ & & \ddots & \alpha_2 \\ O & & & \alpha_1 \end{pmatrix} \cdot \begin{pmatrix} \varphi(a_1) & \varphi(a_2) & \cdots & \varphi(a_n) \\ & \varphi(a_1) & \ddots & \vdots \\ & & \ddots & \varphi(a_2) \\ & & & \varphi(a_1) \end{pmatrix} \cdot \begin{pmatrix} \alpha'_1 & \alpha'_2 & \cdots & \alpha'_n \\ & \alpha'_1 & \ddots & \vdots \\ & & \ddots & \alpha'_2 \\ & & & \alpha'_1 \end{pmatrix}\\
&\equiv (\alpha_1,...,\alpha_n) \cdot E\left((a_1,...,a_n)\right) \cdot (\alpha'_1,...,\alpha'_n).
\end{aligned}
$$

By (i) and (ii), we can conclude that E is a $B = \mathcal{C}^n$-functional. ∎

By the previous lemma, we have a new NCPSpace over $B = \mathcal{C}^n$, (T, E), in the sense of Section 1.1. On this framework, we can do the amalgamated R-transform Calculus, like in Section 1.2.

Lemma 2.2. *Fix $n \in \mathbb{N}$. Let $B = \mathcal{C}^n$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace. Let (T, E) be a Toeplitz matricial probability space over B, induced by (A, φ). Then $C_T(B) = B$, where $C_T(B)$ is the centralizer of B in T, i.e*

$$C_T(B) = \{b \in B : bt = ta, \textit{ for all } t \in T\}.$$

Proof. Let $(\alpha_1, ..., \alpha_n) \in \mathcal{C}^n = B$. Then $(\alpha_1, ..., \alpha_n) \in C_T(B)$. i.e

$$
\begin{aligned}
&(\alpha_1,...,\alpha_n) \cdot (a_1,...,a_n)\\
&\quad = \left(\alpha_1 a_1, \textstyle\sum_{k=1}^{2} \alpha_k a_{(2+1)-k}, ..., \sum_{k=1}^{n} \alpha_k a_{(n+1)-k}\right)\\
&\quad = \left(a_1 \alpha_1, \textstyle\sum_{k=1}^{2} a_{(2+1)-k}\alpha_k, ..., \sum_{k=1}^{n} a_{(n+1)-k}\alpha_k\right)\\
&\quad = \left(a_1 \alpha_1, \textstyle\sum_{j=1}^{2} a_j \alpha_{(2+1)-j}, ..., \sum_{j=1}^{n} a_j \alpha_{(n+1)-j}\right)\\
&\quad = (a_1,...,a_n) \cdot (\alpha_1,...,\alpha_n),
\end{aligned}
$$

for all $(a_1, ..., a_n) \in T$.

Trivially, $C_T(B) \subseteq B$, by the very definition of $C_T(B)$. Therefore,

$$C_T(B) = B.$$

∎

Fix $N \in \mathbb{N}$. Like in Section 1.3, since $C_T(B) = B$, trivial R-transforms of $B = \mathcal{C}^N$-valued random variables contain all information of operator-valued distributions of those B-valued random variables (Also see [32] and [33]). Hence, it is sufficient to consider trivial R-transforms of B-valued random variables to study operator valued random variables, in $\sum_B^s$, where $B = \mathcal{C}^N$ and $s \in \mathbb{N}$. Remark that to study such trivial R-transforms, we need to define suitable cumulants $(K_n)_{n=1}^{\infty}$, as a cumulant multiplicative bimodule map induced by a operator-valued (moment multiplicative) conditional expectation E. Recall that

$$E\left((a_1, ..., a_n)\right) \overset{def}{=} (\varphi(a_1), ..., \varphi(a_n)) \in \mathcal{C}^n,$$

for all $(a_1, ..., a_N) \in T$. Denote cumulants induced by φ, with respect to (A, φ), by $(k_n)_{n=1}^{\infty}$. i,e if $x_1, ..., x_s \in (A, \varphi)$ are random variables, then

$$k_n\left(x_{i_1}, ..., x_{i_n}\right) \overset{def}{=} \sum_{\pi \in NC(n)} \widehat{\varphi}(\pi)\left(x_{i_1} \otimes ..... \otimes x_{i_n}\right) \mu(\pi, 1_n) \in \mathbb{C},$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$, by the Möbius inversion.

Suppose that $(a_1^{(1)}, ..., a_N^{(1)})$,...., $(a_1^{(s)}, ..., a_N^{(s)}) \in T_N$, for the fixed $N \in \mathbb{N}$. Observe the following relation, for $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$;

$$(a_1^{(i_1)}, ..., a_N^{(i_1)})(a_1^{(i_2)}, ..., a_N^{(i_2)}) = (a_1^{(i_1)}a_1^{(i_2)}, \textstyle\sum_{k=1}^{2} a_k^{(i_1)} a_{(2+1)-k}^{(i_2)}, ..., \sum_{k=1}^{N} a_k^{(i_1)} a_{(N+1)-k}^{(i_2)}).$$

Put

$$P_j^{(i_1,i_2)} = \textstyle\sum_{k=1}^{j} a_k^{(i_1)} a_{(j+1)-k}^{(i_2)} \in (A, \varphi) \qquad (\ j = 1, ..., N).$$

i.e

$$(a_1^{(i_1)}a_1^{(i_2)}, \textstyle\sum_{k=1}^{2} a_k^{(i_1)} a_{(2+1)-k}^{(i_2)}, ..., \sum_{k=1}^{N} a_k^{(i_1)} a_{(N+1)-k}^{(i_2)}) = \left(P_1^{(i_1,i_2)}, ..., P_N^{(i_1,i_2)}\right).$$

Then

$$\begin{aligned}&(a_1^{(i_1)}, ..., a_N^{(i_1)})(a_1^{(i_2)}, ..., a_N^{(i_2)})(a_1^{(i_3)}, ..., a_N^{(i_3)})\\ &\quad = \left(P_1^{(i_1,i_2)}, ..., P_N^{(i_1,i_2)}\right)(a_1^{(i_3)}, ..., a_N^{(i_3)})\\ &\quad = \left(P_1^{(i_1,i_2)}a_1^{(i_3)}, \textstyle\sum_{k=1}^{2} P_k^{(i_1,i_2)} a_{(2+1)-k}^{(i_3)}, ..., \sum_{k=1}^{N} P_N^{(i_1,i_2)} a_{(N+1)-k}^{(i_3)}\right).\end{aligned}$$

Similarly, put

$$P_j^{(i_1,i_2,i_3)} = \textstyle\sum_{k=1}^{j} P_k^{(i_1,i_2)} a_{(j+1)-k}^{(i_3)} \qquad (j = 1, ..., N).$$

i.e

$$\begin{aligned}&\left(P_1^{(i_1,i_2)}a_1^{(i_3)}, \textstyle\sum_{k=1}^{2} P_k^{(i_1,i_2)} a_{(2+1)-k}^{(i_3)}, ..., \sum_{k=1}^{N} P_N^{(i_1,i_2)} a_{(N+1)-k}^{(i_3)}\right)\\ &\qquad\qquad = \left(P_1^{(i_1,i_2,i_3)}, ..., P_N^{(i_1,i_2,i_3)}\right).\end{aligned}$$

Inductively, we can define

$$P_j^{(i_1,...,i_{n-1},i_n)} = \sum_{k=1}^{j} P_k^{(i_1,...,i_{n-1})} a_{(j+1)-k}^{(i_n)} \quad (j = 1, ..., N).$$

i.e

$$(a_1^{(i_1)}, ..., a_N^{(i_1)}) \cdots (a_1^{(i_n)}, ..., a_N^{(i_n)}) = \left(P_1^{(i_1,..,i_n)}, ..., P_N^{(i_1,...,i_n)}\right).$$

Definition 2.2. *Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace with its linear functional $\varphi : A \to \mathbb{C}$. Let $(T_N, E_N) \equiv (T, E)$ be a Toeplitz matricial probability space over B with its B-functional $E : T \to B$ and let $(a_1^{(1)}, ..., a_N^{(1)}), ..., (a_1^{(s)}, ..., a_N^{(s)}) \in (T, E)$ be B-valued random variables ($s \in \mathbb{N}$).*

(1) For $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$, define

$$P_j^{(i_1)} = a_j^{(i_1)} \in (A, \varphi), \quad j = 1, ..., N,$$

$$P_j^{(i_1,i_2)} = \sum_{k=1}^{j} a_k^{(i_1)} a_{(j+1)-k}^{(i_2)}, \quad j = 1, ..., N$$

and

$$P_j^{(i_1,...,i_{n-1},i_n)} = \sum_{k=1}^{j} P_k^{(i_1,...,i_{n-1})} a_{(j+1)-k}^{(i_n)}, \quad j = 1, ..., N.$$

Then

$$(a_1^{(i_1)}, ..., a_N^{(i_1)}) \cdots (a_1^{(i_n)}, ..., a_N^{(i_n)}) = \left(P_1^{(i_1,...,i_n)}, ..., P_N^{(i_1,...,i_n)}\right).$$

(2) Define a $(i_1, ..., i_n)$-th moment of $(a_1^{(1)}, ..., a_N^{(1)}), ..., (a_1^{(s)}, ..., a_N^{(s)})$ by

$$E\left((a_1^{(i_1)}, ..., a_N^{(i_1)}) \cdots (a_1^{(i_n)}, ..., a_N^{(i_n)})\right) = \left(\varphi(P_1^{(i_1,...,i_n)}), ..., \varphi(P_N^{(i_1,...,i_n)})\right),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$.

Notice that

$$\varphi\left(P_1^{(i_1,...,i_n)}\right) = \varphi\left(a_1^{(i_1)} ... a_1^{(i_n)}\right)$$

is the $(i_1, ..., i_n)$-th moment of $a_1^{(1)}, ..., a_1^{(s)}$. As we have seen before (and in [32] and [33]), all **trivial** moments contain all information about moments, since $C_T(B) = B$. Hence WLOG, we can only consider the trivial moments of $(a_1^{(1)}, ..., a_N^{(1)})$, ..., $(a_1^{(s)}, ..., a_N^{(s)})$.

Example 2.1. *Suppose that* $N = 2$ *and fix* $B = \mathcal{C}^2$. *Let* (T_2, E_2) *be a Toeplitz matricial probability space over* B, *induced by a NCPSpace* (A, φ). *Let* $(a_1^{(1)}, a_2^{(1)})$, $(a_1^{(2)}, a_2^{(2)})$, $(a_1^{(3)}, a_2^{(3)}) \in (T_2, E_2)$ *be* B*-valued random variables. Then*

$$E\left((a_1^{(1)}, a_2^{(1)}) \cdot (a_1^{(2)}, a_2^{(2)}) \cdot (a_1^{(3)}, a_2^{(3)})\right)$$

$$\equiv E\left(\begin{pmatrix} a_1^{(1)} & a_2^{(1)} \\ O & a_1^{(1)} \end{pmatrix}\begin{pmatrix} a_1^{(2)} & a_2^{(2)} \\ O & a_1^{(2)} \end{pmatrix}\begin{pmatrix} a_1^{(3)} & a_2^{(3)} \\ O & a_1^{(3)} \end{pmatrix}\right)$$

where $O = 0_B = 0_A \in A$.

$$= E\left(\begin{pmatrix} a_1^{(1)}a_1^{(2)}a_1^{(3)} & a_1^{(1)}a_1^{(2)}a_2^{(3)} + a_1^{(1)}a_2^{(2)}a_1^{(3)} + a_2^{(1)}a_1^{(2)}a_1^{(3)} \\ & \\ O & a_1^{(1)}a_1^{(2)}a_1^{(3)} \end{pmatrix}\right)$$

$$\equiv E\left((a_1^{(1)}a_1^{(2)}a_1^{(3)},\ \ a_1^{(1)}a_1^{(2)}a_2^{(3)} + a_1^{(1)}a_2^{(2)}a_1^{(3)} + a_2^{(1)}a_1^{(2)}a_1^{(3)})\right)$$

$$= \left(\varphi(a_1^{(1)}a_1^{(2)}a_1^{(3)}),\ \ \varphi(a_1^{(1)}a_1^{(2)}a_2^{(3)} + a_1^{(1)}a_2^{(2)}a_1^{(3)} + a_2^{(1)}a_1^{(2)}a_1^{(3)})\right).$$

We have that

$$P_1^{(1,2,3)} = a_1^{(1)}a_1^{(2)}a_1^{(3)}$$

and

$$P_2^{(1,2,3)} = a_1^{(1)}a_1^{(2)}a_2^{(3)} + a_1^{(1)}a_2^{(2)}a_1^{(3)} + a_2^{(1)}a_1^{(2)}a_1^{(3)}.$$

2.2. **Toeplitz Matricial Cumulants.**

Throughout this section, we will fix $N \in \mathbb{N}$. Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace, with its linear functional $\varphi : A \to \mathbb{C}$. Let $(T_N, E_N) \equiv (T, E)$ be a Toeplitz matricial probability space over B with its B-functional $E : T \to B$,

$$E\left((a_1, ..., a_N)\right) = (\varphi(a_1), ..., \varphi(a_N)) \in B,$$

for all $(a_1, ..., a_N) \in T$.Let $(a_1^{(1)}, ..., a_N^{(1)}), ..., (a_1^{(s)}, ..., a_N^{(s)}) \in (T, E)$ be B-valued random variables ($s \in \mathbb{N}$). Then we can conpute the $(i_1, ..., i_n)$-th moment of them by

$$\begin{aligned} &E\left((a_1^{(i_1)}, ..., a_N^{(i_1)})...(a_1^{(i_n)}, ..., a_N^{(i_n)})\right) \\ &\qquad = E\left((P_1^{(i_1,...,i_n)}, ..., P_N^{(i_1,...,i_n)})\right) \\ &\qquad = \left(\varphi(P_1^{(i_1,...,i_n)}), ..., \varphi(P_N^{(i_1,...,i_n)})\right), \end{aligned}$$

where

$$P_j^{(i_1)} = a_j^{(i_1)} \in (A, \varphi),$$

$$P_j^{(i_1,i_2)} = \textstyle\sum_{k=1}^{j} a_k^{(i_1)} a_{(j+1)-k}^{(i_2)} \in (A, \varphi),$$

and

$$P_j^{(i_1,...,i_{n-1},i_n)} = \textstyle\sum_{k=1}^{j} P_k^{(i_1,...,i_{n-1})} a_{(j+1)-k}^{(i_n)} \in (A, \varphi),$$

for all $j = 1, ..., N$.

So, we can define a moment series of $A_1 = (a_1^{(1)}, ..., a_N^{(1)}), ..., A_s = (a_1^{(s)}, ..., a_N^{(s)})$ denoted by $M_{A_1,...,A_s}$, by

$$\begin{aligned} &M_{A_1,...,A_s}(z_1, ..., z_s) = M^t_{A_1,...,A_s}(z_1, ..., z_s) \\ &\quad = \textstyle\sum_{n=1}^{\infty} \sum\limits_{i_1,...,i_n \in \{1,...,s\}} E\left(A_{i_1}...A_{i_n}\right) z_{i_1}...z_{i_n} \\ &\quad = \textstyle\sum_{n=1}^{\infty} \sum\limits_{i_1,...,i_n \in \{1,...,s\}} \left(\varphi(P_1^{(i_1,...,i_n)}), ..., \varphi(P_N^{(i_1,...,i_n)})\right) z_{i_1}...z_{i_n}. \end{aligned}$$

In this section, we will show that

$$\begin{aligned} &K_n\left((a_1^{(i_1)}, ..., a_N^{(i_1)}), ..., (a_1^{(i_n)}, ..., a_N^{(i_n)})\right) \\ &\qquad\qquad = \left(k_n(Q_1^{(i_1,...,i_n)}), ..., k_n(Q_N^{(i_1,...,i_n)})\right), \end{aligned}$$

where $(k_n)_{n=1}^{\infty}$ is the cumulants induced by φ, in the sense of Nica and Speicher, and

$$Q_j^{(i_1)} = a_j^{(i_1)}, \;\; j = 1, ..., N,$$

$$Q_j^{(i_1,i_2)} = \textstyle\sum_{k=1}^{2} \left(a_k^{(i_1)}, a_{(j+1)-k}^{(i_2)}\right) \in A \times A, \;\; j = 1, ..., N$$

and

$$Q_j^{(i_1,...,i_{n-1},i_n)} = \sum_{k=1}^{n} \left(Q_j^{(i_1,...,i_{n-1})}, a_{(j+1)-k}^{(i_n)} \right) \in \underbrace{A \times .... \times A}_{n-times},$$

for all $j = 1, ..., N$. Actually, we can regard $\underbrace{A \times ... \times A}_{n-times}$ as $\underbrace{A \otimes_B .... \otimes_B A}_{n-times}$, since $C_T(B) = B$, whenever $B = \mathcal{C}^N$ and T is a Toeplitz matricial probability space over B. So,

$$\begin{aligned} & k_n \left(\sum_{k=1}^{n} \left(Q_j^{(i_1,...,i_{n-1})}, a_{(j+1)-k}^{(i_n)} \right) \right) \\ & \quad = \sum_{k=1}^{n} k_n \left(Q_j^{(i_1,...,i_{n-1})}, a_{(j+1)-k}^{(i_n)} \right) \\ & \quad = \sum_{k=1}^{n} c^{(n)} \left(Q_j^{\otimes(i_1,...,i_{n-1})} \otimes a_{(j+1)-k}^{(i_n)} \right), \end{aligned}$$

is well-detemined, where $c^{(n)}$ is the n-thcomponent of the cumulant multiplicative function $\widehat{c} = (c^{(k)})_{k=1}^{\infty} \in I(A, \mathbb{C})$, induced by φ and where

$$Q_j^{\otimes(i_1,...,i_{n-1})} = Q_j^{\otimes(i_1,...,i_{n-2})} \otimes a_{(j+1)-k}^{(i_{n-1})},$$

with

$$Q_j^{\otimes(i_1)} = a_j^{(i_1)} \quad \text{and} \quad Q_j^{\otimes(i_1,i_2)} = \sum_{k=1}^{2} a_k^{(i_1)} \otimes a_{(j+1)-k}^{(i_2)},$$

for all $j = 1, ..., N$.

Definition 2.3. *Fix $s \in \mathbb{N}$. Let*

$$P_j^{(i_1,...,i_n)} = \sum_{k=1}^{j} P_k^{(i_1,...,i_{n-1})} a_{(j+1)-k}^{(i_n)} \in (A, \varphi)$$

with

$$P_j^{(i_1)} = a_j^{(i_1)} \quad \textit{and} \quad P_j^{(i_1,i_2)} = \sum_{k=1}^{j} a_k^{(i_1)} a_{(j+1)-k}^{(i_2)},$$

in (A, φ), be determined inductively, for all $j = 1, ..., N$. Then, with respect to them, we will define

$$\begin{aligned} & Q_j^{(i_1)} = a_j^{(i_1)} \in A, \\ & Q_j^{(i_1,i_2)} = \sum_{k=1}^{j} \left(a_j^{(i_1)}, a_{(j+1)-k}^{(i_2)} \right) \in A \times A \end{aligned}$$

and inductively on n,

$$Q_j^{(i_1,...,i_n)} = \sum_{k=1}^{j} \left(Q_j^{(i_1,...,i_{n-1})}, a_{(j+1)-k}^{(i_n)} \right) \in \underbrace{A \times ... \times A}_{n-times},$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$, for all $j = 1, ..., N$.

Example 2.2. *At the end of the previous section, we observed an example such that*

$$P_1^{(1,2,3)} = a_1^{(1)} a_1^{(2)} a_1^{(3)}$$

and

$$P_2^{(1,2,3)} = a_1^{(1)} a_1^{(2)} a_2^{(3)} + a_1^{(1)} a_2^{(2)} a_1^{(3)} + a_2^{(1)} a_1^{(2)} a_1^{(3)}.$$

In this case, we can decide

$$Q_1^{(1,2,3)} = \left(a_1^{(1)}, a_1^{(2)}, a_1^{(3)}\right)$$

and

$$Q_2^{(1,2,3)} = \left(a_1^{(1)}, a_1^{(2)}, a_2^{(3)}\right) + \left(a_1^{(1)}, a_2^{(2)}, a_1^{(3)}\right) + \left(a_2^{(1)}, a_1^{(2)}, a_1^{(3)}\right).$$

Notice that the addition above is meaningless. This addition is not a pointwise addition ! This addition is only depending on cumulant k_n, induced by φ. And we can get that

$$Q_1^{\otimes(1,2,3)} = a_1^{(1)} \otimes a_1^{(2)} \otimes a_1^{(3)}$$

and

$$Q_2^{\otimes(1,2,3)} = \left(a_1^{(1)} \otimes a_1^{(2)} \otimes a_2^{(3)}\right) + \left(a_1^{(1)} \otimes a_2^{(2)} \otimes a_1^{(3)}\right) + \left(a_2^{(1)} \otimes a_1^{(2)} \otimes a_1^{(3)}\right).$$

Proposition 2.3. *Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace. Let (T, E) be a Toeplitz matricial probability space over B and let $A_1 = (a_1^{(1)}, ..., a_N^{(1)}), ..., A_s = (a_1^{(s)}, ..., a_N^{(s)}) \in (T, E)$ be B-valued random variables ($s \in \mathbb{N}$). Then the R-transform of $A_1, ..., A_s$ is determined by*

$$\begin{aligned} R_{A_1,...,A_s}(z_1, ..., z_s) &\overset{def}{=} R^t_{A_1,...,A_s}(z_1, ..., z_s) \\ &= \textstyle\sum_{n=1}^{\infty} \sum\limits_{i_1,...,i_n \in \{1,...,s\}} K_n(A_{i_1}, ..., A_{i_n})\, z_{i_1} ... z_{i_n}, \end{aligned}$$

where

$$K_n(A_{i_1}, ..., A_{i_n}) = \sum_{\pi \in NC(n)} \widehat{E}(\pi)\,(A_{i_1} \otimes ... \otimes A_{i_n})\, \mu(\pi, 1_n),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$.

Proof. By the Möbius inversion, we can always define a $(i_1, ..., i_n)$-th cumulants from moments. We defined that, since $C_T(B) = B$, moments of $A_1, ..., A_s$ by trivial moments of $A_1, ..., A_s$. So, WLOG, we can define a $(i_1, ..., i_n)$-th cumulants of $A_1, ..., A_s$ by trivial (amalgamated) cumulants of them. So, by Section 1.2, we have that

$$\begin{aligned} K_n\left(A_{i_1}, ..., A_{i_n}\right) &= c^{(n)}\left(A_{i_1} \otimes ... \otimes A_{i_n}\right) \\ &= \sum_{\pi \in NC(n)} \widehat{E}(\pi)(A_{i_1} \otimes ... \otimes A_{i_n})\mu(\pi, 1_n), \end{aligned}$$

where $\widehat{E} = (E^{(k)})_{k=1}^{\infty} \in I^m(T, B)$ is the moment multiplicative bimodule map induced by the B-functional $E : T \to B$. Also, again, by Section 1.2, we can define

$$R_{A_1,...,A_s}(z_1, ..., z_s) = R^t_{A_1,...,A_s}(z_1, ..., z_s).$$

∎

Definition 2.4. *Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace with its linear functional $\varphi : A \to \mathbb{C}$. Let $(T_N, E_N) \equiv (T, E)$ be a Toeplitz matricial probability space over B and let $(a_1^{(1)}, ..., a_N^{(1)}), ..., (a_1^{(s)}, ..., a_N^{(s)})$ be B-valued random variables ($s \in \mathbb{N}$). For $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$, define a $(i_1, ..., i_n)$-th cumulants, by*

$$\begin{aligned} &K_n\left((a_1^{(i_1)}, ..., a_N^{(i_1)}), ..., (a_1^{(i_n)}, ..., a_N^{(i_n)})\right) \\ &\quad = \sum_{\pi \in NC(n)} \widehat{E}(\pi)\left((a_1^{(i_1)}, ..., a_N^{(i_1)}) \otimes ... \otimes (a_1^{(i_n)}, ..., a_N^{(i_n)})\right)\mu(\pi, 1_n). \end{aligned}$$

Lemma 2.4. *Let (T, E) be a Toeplitz matricial probability space over $B = \mathcal{C}^N$, induced by (A, φ) and let $A_1 = (a_1^{(1)}, ..., a_N^{(1)})$, ..., $A_s = (a_1^{(s)}, ..., a_N^{(s)}) \in (T, E)$ be $B = \mathcal{C}^N$-valued random varialbes ($s \in \mathbb{N}$). Then $(i_1, ..., i_n)$-th moment of $A_1, ..., A_s$ has the following relation ;*

$$E\left(A_{i_1} ... A_{i_n}\right) = \left(\sum_{\pi \in NC(n)} k_\pi(Q_1^{(i_1,...,i_n)}), ..., \sum_{\pi \in NC(n)} k_\pi\left(Q_N^{(i_1,...,i_n)}\right)\right),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$, where k_π is a cumulants depending on $\pi \in NC(n)$, induced by φ, with respect to (A, φ).

Proof. By the previous observation, we have that the $(i_1, ..., i_n)$-th (trivial) moment of $A_1, ..., A_s$ is

$$\begin{aligned} &E\left(A_{i_1} ... A_{i_n}\right) \\ &\quad = E\left((a_1^{(i_1)}, ..., a_N^{(i_1)}) ... (a_1^{(i_n)}, ..., a_N^{(i_n)})\right) \end{aligned}$$

$$= E\left((P_1^{(i_1,...,i_n)}, ..., P_N^{(i_1,...,i_n)})\right)$$

where

$$P_1^{(i_1,...,i_n)} = a_1^{(i_1)} a_1^{(i_2)} ... a_1^{(i_n)}$$

and

$$P_j^{(i_1,...,i_n)} = \sum_{k=1}^{j} P_k^{(i_1,...,i_{n-1})} a_{(j+1)-k}^{(i_n)}, \text{ for } j = 2, ..., N,$$

hence

$$= \left(\varphi(P_1^{(i_1,...,i_n)}), ..., \varphi(P_N^{(i_1,...,i_n)})\right) \in B.$$

By the Möbius inversion,

$$\begin{aligned}\varphi\left(P_1^{(i_1,...,i_n)}\right) &= \varphi\left(a_1^{(i_1)} a_1^{(i_2)} ... a_1^{(i_n)}\right) \\ &= \sum_{\pi \in NC(n)} k_\pi\left(a_1^{(i_1)}, a_1^{(i_2)}, ..., a_1^{(i_n)}\right)\end{aligned}$$

and

$$\varphi\left(P_j^{(i_1,...,i_n)}\right) = \sum_{\theta \in NC(n)} k_\theta\left(Q_j^{(i_1,...,i_n)}\right),$$

for all $j = 2, ..., N$, by the definition of $Q_j^{(i_1,...,i_n)}$, where above k_π and k_θ, for $\pi, \theta \in NC(n)$, are defined in the sense of Nica and Speicher (See Section 1.3 or see [2] and [3]). Thus

$$\begin{aligned}E\left(A_{i_1} ... A_{i_n}\right) &= \left(\varphi(P_1^{(i_1,...,i_n)}), ..., \varphi(P_N^{(i_1,...,i_n)})\right) \\ &= \left(\sum_{\pi \in NC(n)} k_\pi\left(Q_1^{(i_1,...,i_n)}\right), ..., \sum_{\pi \in NC(n)} k_\pi\left(Q_N^{(i_1,...,i_n)}\right)\right) \in B,\end{aligned}$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. ∎

Theorem 2.5. *Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace with its linear functional $\varphi : A \to \mathbb{C}$. Let (T, E) be a Toeplitz matricial probability space over B with its B-functional $E : T \to B$ and let $A_1 = (a_1^{(1)}, ..., a_N^{(1)}), ..., A_s = (a_1^{(s)}, ..., a_N^{(s)}) \in (T, E)$ be B-valued random variables ($s \in \mathbb{N}$). Then the $(i_1, ..., i_n)$-th cumulant of $A_1, ..., A_s$ is*

$$K_n\left(A_{i_1}, ..., A_{i_n}\right) = \left(k_n(Q_1^{(i_1,...,i_n)}), ..., k_n(Q_N^{(i_1,...,i_n)})\right) \in B$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$.

Proof. By definition, we have that

$$K_n\left(A_{i_1},...,A_{i_n}\right)=\sum_{\pi\in NC(n)}\widehat{E}(\pi)\left(A_{i_1}\otimes...\otimes A_{i_n}\right)\mu(\pi,1_n),$$

for any $(i_1,...,i_n)\in\{1,...,s\}^n$, $n\in\mathbb{N}$. Fix $(i_1,...,i_n)$. Then the right-hand side of the previous formular can be rewritten

$$\sum_{\pi\in NC(n)}\prod_{V\in\pi(o)}\widehat{E}(\pi\mid_V)\left(A_{i_1}\otimes...\otimes A_{i_n}\right)\mu\left(\pi\mid_V,1_n\mid_V\right).$$

Recall that if $B=\mathcal{C}^N$, then $C_T(B)=B$. Thus, for any $(j_1,...,j_k)\in\{1,...,s\}^k$, $k\in\mathbb{N}$,

$$E\left(A_{j_1}\otimes...\otimes A_{j_k}\right)\in B=C_T(B).$$

By the previous property, also we can rewrite the right-hand side of the previous formular, by

$$\sum_{\pi\in NC(n)}\prod_{W\in\pi=\pi(o)\cup\pi(i)}\widehat{E}(\pi\mid_W)\left(A_{i_1}\otimes...\otimes A_{i_n}\right)\mu(\pi\mid_W,1_n\mid_W).\text{——(*)}$$

Now, let $W=(j_1,...,j_k)\in\pi=\pi(o)\cup\pi(i)$ be a block, where $\pi\in NC(n)$. Then, by the previous lemma, we have that

$$\begin{aligned}&\widehat{E}(\pi\mid_W)\left(A_{i_1}\otimes...\otimes A_{i_n}\right)=E^{(k)}\left(A_{j_1}\otimes...\otimes A_{j_k}\right)\\&\quad=E\left(A_{j_1}...A_{j_k}\right)\\&\quad=E\left((a_1^{(j_1)},...,a_N^{(j_1)})...(a_1^{(j_k)},....,a_N^{(j_k)})\right)\\&\quad=E\left((P_1^{(j_1,...,j_k)},...,P_N^{(j_1,...,j_k)})\right)\\&\quad=\left(\varphi(P_1^{(j_1,...,j_k)}),....,\varphi(P_N^{(j_1,...,j_k)})\right)\\&\quad=\left(\sum_{\theta\in NC(k)}k_\theta(Q_1^{(j_1,...,j_k)}),....,\sum_{\theta\in NC(k)}k_\theta(Q_N^{(j_1,...,j_k)})\right)\in B.\end{aligned}$$

Therefore,

$$(*)=\sum_{\pi\in NC(n)}\prod_{W\in\pi}\left(\sum_{\theta\in NC(k)}k_\theta(Q_1^{(W)}),....,\sum_{\theta\in NC(k)}k_\theta(Q_N^{(W)})\right)\mu(\pi\mid_W,1_n\mid_W),$$

where $Q_j^{(W)} = Q_j^{(j_1,...,j_k)}$, if $W = (j_1, ..., j_k) \in \pi \in NC(n)$ $(k < n)$. Equivalently, we have that

$$K_n\left(A_{i_1}, ..., A_{i_n}\right) = \left(k_n(Q_1^{(i_1,...,i_n)}), ..., k_n(Q_N^{(i_1,...,i_n)})\right).$$

▮

The proof of the above theorem is motivated by cumulants of type B studied in [31] (See Section 6.2 and 6.3 in [31]).

Example 2.3. *We will use the same example which we have observed. Suppose we are given* $A_1 = (a_1^{(1)}, a_2^{(1)})$, $A_2 = (a_1^{(2)}, a_2^{(2)})$ *and* $A_3 = (a_1^{(3)}, a_2^{(3)})$, $\mathcal{C}^2$*-valued random variables in* (T_2, E_2). *Then we have that*

$$Q_1^{(1,2,3)} = \left(a_1^{(1)}, a_1^{(2)}, a_1^{(3)}\right)$$

and

$$Q_2^{(1,2,3)} = \left(a_1^{(1)}, a_1^{(2)}, a_2^{(3)}\right) + \left(a_1^{(1)}, a_2^{(2)}, a_1^{(3)}\right) + \left(a_2^{(1)}, a_1^{(2)}, a_1^{(3)}\right).$$

So,

$$K_3\left(A_1, A_2, A_3\right) = \left(k_3(Q_1^{(1,2,3)}),\ k_3(Q_2^{(1,2,3)})\right)$$

$$= (k_3(a_1^{(1)}, a_1^{(2)}, a_1^{(3)}),$$
$$k_3\left(a_1^{(1)}, a_1^{(2)}, a_2^{(3)}\right) + k_3\left(a_1^{(1)}, a_2^{(2)}, a_1^{(3)}\right) + k_3\left(a_2^{(1)}, a_1^{(2)}, a_1^{(3)}\right)).$$

2.3. **Toeplitz Matricial R-transform Theory.**

Let (A, φ) be a NCPSpace, in the sense of Section 1.3, with its linear functional $\varphi : A \to \mathbb{C}$. In this section, by using information about cumulants, we will consider R-transform Calculus on a Toeplitz matricial probability space (T, E), over a Toeplitz matricial algebra, $B = \mathcal{C}^N$, induced by a NCPSpace (A, φ). Also, we will observe the freeness of subsets of T. To do that, we need to observe the freeness on (T, E). Clearly, by the Speicher's characterization, we can define freeness on (T, E). We want to express this freeness with respect to the freeness on (A, φ).

Let $X_1 = (x_1^{(1)}, ..., x_N^{(1)}), ..., X_s = (x_1^{(s)}, ..., x_N^{(s)})$, $Y_1 = (y_1^{(1)}, ..., y_N^{(1)}), ...,$ and $Y_t = (y_1^{(t)}, ..., y_N^{(t)})$ be $B = \mathcal{C}^N$-valued random variables ($s, t \in \mathbb{N}$). Define two subsets of T,

$$\Xi = \{X_1, ..., X_s\} \text{ and } \Upsilon = \{Y_1, ..., Y_t\}.$$

By Speicher's (amalgamated) Freeness characterization, we can say that

[Two subsets Ξ and Υ are free over B]

$\Longleftrightarrow$[All mixed B-valued cumulants vanish].

Remark that, since $C_T(B) = B$, "all mixed cumulants" can be replaced by "all **trivial** mixed cumulants" and these trivial cumulants is defined by $B = \mathcal{C}^N$-valued cumulants, above. Also, remark that, since $C_T(B) = B$, we can avoid the insertion property when we compute the B-valued cumulants. By the previous section, if $A_{i_1}, ..., A_{i_n} \in (T, E)$ are B-valued random variables, then we have that

$$\begin{aligned}
&K_n\left(A_{i_1}, ..., A_{i_n}\right) \overset{def}{=} \sum_{\pi \in NC(n)} \widehat{E}(\pi)\left(A_{i_1} \otimes ... \otimes A_{i_n}\right) \mu(\pi, 1_n) \\
&= \sum_{\pi \in NC(n)} \prod_{V \in \pi} \widehat{E}(\pi \mid_V)\left(A_{i_1} \otimes ... \otimes A_{i_n}\right) \mu(\pi \mid_V, 1_{|V|}) \\
&= \sum_{\pi \in NC(n)} \prod_{V \in \pi} \left(\varphi(P_1^{(i_1, ..., i_n)}), ..., \varphi(P_N^{(i_1, ..., i_n)})\right) \mu(\pi \mid_V, 1_{|V|}) \\
&= \sum_{\pi \in NC(n)} \prod_{V \in \pi} \left(\sum_{\theta \in NC(|V|)} k_\theta(Q_1^{(V)}), ..., \sum_{\theta \in NC(|V|)} k_\theta(Q_N^{(V)})\right) \mu(\pi \mid_V, 1_{|V|}) \\
&= \left(k_n(Q_1^{(i_1, ..., i_n)}), ..., k_n(Q_N^{(i_1, ..., i_n)})\right) \in B,
\end{aligned}$$

where K_n's are B-valued cumulants induced by E and k_n's are cumulants, in the sense of Nica and Speicher, induced by φ and where

$$P_j^{(i_1, ..., i_n)} = \sum_{k=1}^{j} P_k^{(i_1, ..., i_{n-1})} a_{(j+1)-k}^{(i_n)} \in (A, \varphi),$$

and

$$Q_j^{(i_1, ..., i_n)} = \sum_{k=1}^{j} \left(Q_k^{(i_1, ..., i_{n-1})}, a_{(j+1)-k}^{(i_n)}\right) \in \underbrace{A \times ... \times A}_{n-times},$$

for all $j = 1, ..., N$.

Theorem 2.6. *Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (A, φ) be an arbitrary NCPSpace. Let $(T_N, E_N) \equiv (T, E)$ be a Toeplitz matricial probability space over B and let $X_1 = (x_1^{(1)}, ..., x_N^{(1)}), ..., X_s = (x_1^{(s)}, ..., x_N^{(s)}), Y_1 = (y_1^{(1)}, ..., y_N^{(1)}), ...,$ and $Y_t = (y_1^{(t)}, ..., y_N^{(t)})$ be B-valued random variables ($s, t \in \mathbb{N}$). Define subsets $\Xi = \{X_1, ..., X_s\}$ and $\Upsilon = \{Y_1, ..., Y_t\}$ in T. Then Ξ and Υ are free over B, in (T, E) if and only if*

Proof. Suppose that Ξ and Υ are free over B, in (T, E). Then, by definition, all B-valued mixed cumulants of Ξ and Υ vanish ;

$$K_n\left(A_{i_1}, ..., A_{i_n}\right) = \left(k_n\left(Q_1^{(i_1,...,i_n)}\right), ..., k_n\left(Q_N^{(i_1,...,i_n)}\right)\right) = 0_B,$$

where

$$0_B \equiv \begin{pmatrix} 0 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & 0 \end{pmatrix}_{N\times N} \equiv (0, ..., 0) \in B = \mathcal{C}^N$$

and $A_{i_1}, ..., A_{i_n} \in \Xi \cup \Upsilon$ (mixed) in (T, E). It suffices to consider the case when $A_{i_j} = Y_{i_j}$ and $A_{i_k} = X_{i_k}$, $k \neq j$ in $\{1, ..., n\}$. i.e put

$$A_{i_1} = X_{i_1}, ..., A_{i_{j-1}} = X_{i_{j-1}}, \boxed{A_{i_j} = Y_{i_j}}, A_{i_{j+1}} = X_{i_{j+1}} ..., A_{i_n} = X_{i_n}.$$

First, we will consider $P_1^{(i_1,...,i_n)}$,..., $P_N^{(i_1,...,i_n)}$, because $Q_j^{(i_1,...,i_n)}$'s are determined by them ;

$$P_1^{(i_1,...,i_n)} = x_1^{(i_1)} ... x_1^{(i_{j-1})} y_1^{(i_j)} x_1^{(i_{j+1})} ... x_1^{(i_n)}$$

and

$$P_j^{(i_1,...,i_n)} = \textstyle\sum_{k=1}^{j} P_j^{(i_1,...,i_{n-1})} a_{(j+1)-k}^{(i_n)}, \text{ for all } j = 1, ..., N.$$

Note that if $i_j = i_n$, then $P_j^{(i_1,...,i_{n-1},i_n)}$ has the form of sum of mixed words in $x_1^{(i_1)}, ..., x_N^{(i_1)}, ..., x_1^{(i_{n-1})}, ..., x_N^{(i_{n-1})}$ and $\boxed{y_1^{(i_j)} = y_1^{(i_n)}, ..., y_N^{(i_j)} = y_N^{(i_n)}}$. If $1 < j < n$, then, also, $P_j^{(i_1,...,i_j,...,i_n)}$ has the form of sum of mixed words in $x_1^{(i_1)}, ..., x_N^{(i_1)}, ..., x_1^{(i_{j-1})}, ..., x_N^{(i_{j-1})}$, $\boxed{y_1^{(i_j)}, ..., y_N^{(i_j)}}$, $x_1^{(i_{j+1})}, ..., x_N^{(i_{j+1})}$, ..., $x_1^{(i_n)}, ..., x_N^{(i_n)}$. We can conclude, by induction. So, under our assumption, generally, for $1 \leq j \leq n$,

$$\begin{aligned} K_n\left(A_{i_1}, ..., A_{i_n}\right) &= \left(k_n\left(Q_1^{(i_1,...,i_n)}\right), ..., k_n\left(Q_N^{(i_1,...,i_n)}\right)\right) \\ &= (k_n(x_1^{(i_1)}, ..., y_1^{(i_j)}, ..., x_1^{(i_n)}), \textstyle\sum_{k=1}^{2} k_n(Q_2^{(i_1,...,i_{n-1})}, a_{(2+1)-k}^{(i_n)}), \\ &\qquad ..., \textstyle\sum_{k=1}^{N} k_n(Q_N^{(i_1,...,i_{n-1})}, a_{(j+1)-k}^{(i_n)})) = (0, ..., 0) \in B. \end{aligned}$$

$\Longleftrightarrow$

$k_n(x_1^{(i_1)}, ..., x_1^{(i_{j-1})}, y_1^{(i_j)}, x_1^{(i_{j+1})}, ..., x_1^{(i_n)}) = 0 \in \mathbb{C}$,

$\sum_{k=1}^{2} k_n(Q_2^{(i_1,...,i_{n-1})}, a_{(2+1)-k}^{(i_n)}) = 0 \in \mathbb{C}$,

$$\vdots$$

$\sum_{k=1}^{N} k_n(Q_N^{(i_1,...,i_{n-1})}, a_{(j+1)-k}^{(i_n)}) = 0 \in \mathbb{C}.$

$\Longleftrightarrow$

All mixed cumulants of $\{x_1^{(i_1)}, ..., x_N^{(i_1)}, ..., x_1^{(i_{j-1})}, ..., x_N^{(i_{j-1})}, x_1^{(i_{j+1})}, ..., x_N^{(i_{j+1})}, ..., x_1^{(i_n)}, ..., x_N^{(i_n)}\}$ and $\{y_1^{(i_j)}, ..., y_N^{(i_j)}\}$ are free (over $\mathbb{C}$), in the sense of Section 1.3, in (A, φ). Therefore, we can get that

$\Xi = \{X_1 = (x_1^{(1)}, ..., x_N^{(1)}), ..., X_s = (x_1^{(s)}, ..., x_N^{(s)})\}$ and $\Upsilon = \{Y_1 = (y_1^{(1)}, ..., y_N^{(1)}), ..., Y_t = (y_1^{(t)}, ..., y_N^{(t)})\}$ are free over $B = \mathcal{C}^N$, in (T, E) if and only if $\{x_1^{(l)}, ..., x_N^{(l)} : l = 1, ..., s\}$ and $\{y_1^{(l')}, ..., y_N^{(l')} : l' = 1, ..., t\}$ are free, in the sense of Section 1.3, in (A, φ). ∎

Let

$$X_1 \equiv \begin{pmatrix} x_1^{(1)} & x_2^{(1)} & \cdots & x_N^{(1)} \\ & x_1^{(1)} & \ddots & \vdots \\ & & \ddots & x_2^{(1)} \\ O & & & x_1^{(1)} \end{pmatrix}, ..., X_s \equiv \begin{pmatrix} x_1^{(s)} & x_2^{(s)} & \cdots & x_N^{(s)} \\ & x_1^{(s)} & \ddots & \vdots \\ & & \ddots & x_2^{(s)} \\ O & & & x_1^{(s)} \end{pmatrix}$$

and

$$Y_1 \equiv \begin{pmatrix} y_1^{(1)} & y_2^{(1)} & \cdots & y_N^{(1)} \\ & y_1^{(1)} & \ddots & \vdots \\ & & \ddots & y_2^{(1)} \\ O & & & y_1^{(1)} \end{pmatrix}, ..., Y_t = \begin{pmatrix} y_1^{(t)} & y_2^{(t)} & \cdots & y_N^{(t)} \\ & y_1^{(t)} & \ddots & \vdots \\ & & \ddots & y_2^{(t)} \\ O & & & y_1^{(t)} \end{pmatrix}$$

be $\mathcal{C}^N$-valued random variables in (T_N, E_N), induced by (A, φ). Then, by the previous theorem, we characterize $\mathcal{C}^N$-valued freeness with respect to (scarlar-valued) freeness on (A, φ). i.e $\{X_1, ..., X_s\}$ and $\{Y_1, ..., Y_t\}$ are free over $\mathcal{C}^N$ if and only if $\{x_1^{(l)}, ..., x_N^{(l)} : l = 1, ..., s\}$ and $\{y_1^{(l')}, ..., y_N^{(l')} : l' = 1, ..., t\}$ are free in (A, φ).

Now, we can characterize $\mathcal{C}^N$-valued freeness on (T_N, E_N) with respect to scalar-valued freeness on (A, φ), where (T_N, E_N) is a Toeplitz matricial probability space over a Toeplitz matricial algebra, induced by (A, φ). So, we can observe R-transform Calculus.

Proposition 2.7. *Let (A, φ) be a NCPSpace and let $x_1^{(l)}, ..., x_N^{(l)}, y_1^{(l')}, ..., y_N^{(l')} \in (A, \varphi)$ be random variables, where $l = 1, ..., s$ and $l' = 1, ..., t$ $(s, t \in \mathbb{N})$. Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (T, E) be a NCPSpace over B, induced by (A, φ). Let*

$$X_l = (x_1^{(l)}, ..., x_N^{(l)}) \quad \textit{and} \quad Y_{l'} = (y_1^{(l')}, ..., y_N^{(l')})$$

be B-valued random variables in (T, E), where $l = 1, ..., s$ and $l' = 1, ..., t$. If $\widetilde{X} = \{x_1^{(l)}, ..., x_N^{(l)} : l = 1, ..., s\}$ and $\widetilde{Y} = \{y_1^{(l')}, ..., y_N^{(l')} : l' = 1, ..., t\}$ are free in (A, φ), then

(1) $R_{X_1,...,X_s,Y_1,...,Y_t}(z_1, ..., z_{s+t}) = R_{X_1,...,X_s}(z_1, ..., z_s) + R_{Y_1,...,Y_t}(z_{s+1}, ..., z_{s+t})$

(2) if $s = t$, then

$$R_{X_1+Y_1,...,X_s+Y_s}(z_1, ..., z_s) = R_{X_1,...,X_s}(z_1, ..., z_s) + R_{Y_1,...,Y_s}(z_1, ..., z_s)$$

(3) if $s = t$, then

$$R_{X_1Y_1,...,X_sY_s}(z_1, ..., z_s) = (R_{X_1,...,X_s} \boxed{*} R_{Y_1,...,Y_s})(z_1, ..., z_s).$$

Proof. By the previous theorem, if $\widetilde{X}$ and $\widetilde{Y}$ are free in (A, φ), then $\{X_1, ..., X_s\}$ and $\{Y_1, ..., Y_t\}$ are free over B, in (T, E). So, (1) and (2) are easily verified. We will only observe (3), when $s = t$. Fix $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. Then

$$K_n\left(X_{i_1}Y_{i_1}, ..., X_{i_n}Y_{i_n}\right) = \left(k_n(Q_1^{(i_1,...,i_n)}), ..., k_n(Q_N^{(i_1,...,i_n)})\right),$$

where

$$\begin{aligned} X_{i_k}Y_{i_k} &= (x_1^{(i_k)}, ..., x_N^{(i_k)}) \cdot (y_1^{(i_k)}, ..., y_N^{(i_k)}) \\ &= \left(x_1^{(i_k)}y_1^{(i_k)}, \sum_{k=1}^{2} x_k^{(i_1)} y_{(2+1)-k}^{(i_k)}, ..., \sum_{k=1}^{N} x_k^{(i_n)} x_{(N+1)-k}^{(i_n)}\right) \\ &\equiv (a_1^{(i_k)}, ..., a_N^{(i_k)}), \end{aligned}$$

for all $k = 1, ..., n$ and

$$Q_j^{(i_1,...,i_n)} = \sum_{k=1}^{j} \left(Q_j^{(i_1,...,i_{n-1})}, a_{(j+1)-k}^{(i_n)}\right) \in \underbrace{A \times ... \times A}_{n-times},$$

for all $j = 1, ..., N$. Thus,

$$k_n\left(Q_j^{(i_1,...,i_n)}\right) = \sum_{k=1}^{j} k_n\left(Q_j^{(i_1,...,i_{n-1})}, a_{(j+1)-k}^{(i_n)}\right)$$

and, inductively, $\left(Q_j^{(i_1,...,i_{n-1})}, a_{(j+1)-k}^{(i_n)}\right)$ is the sum of alternating tuples of $x_p^{(q)}$'s and $y_{p'}^{(q')}$'s. By little abuse of notation, we have that

$$\left(k_n(Q_1^{(i_1,...,i_n)}), ..., k_n(Q_N^{(i_1,...,i_n)})\right)$$

$$= (\sum_{\theta \in NC(n)} k_{\theta}(Q_1(x_p^{(q)})) \cdot k_{Kr(\theta)}(Q_1(y_{p'}^{(q')})), ...$$

$$..., \sum_{\theta \in NC(n)} k_{\theta}(Q_N(x_p^{(q)}) \cdot k_{Kr(\theta)}(Q_N(y_{p'}^{(q')}))),$$

where $\theta \cup Kr(\theta) \in NC(2n)$ is the alternating union of noncrossing partitions such that $x_p^{(q)}$'s are depending on θ and $y_{p'}^{(q')}$'s are depending on $Kr(\theta)$. Therefore,

$$K_n \left(X_{i_1} Y_{i_1}, ..., X_{i_n} Y_{i_n} \right) = \sum_{\pi \in NC(n)} \left(K_{\pi}(X_{i_1}, ..., X_{i_n}) \right) \left(K_{Kr(\pi)}(Y_{i_1}, ..., Y_{i_n}) \right).$$

(Again, remark that $C_T(B) = B$!!) ∎

3. Applications

3.1. **Operator-Valued Moments and Cumulants of a Single Toeplitz matricial valued random variable.**

Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace, in the sense of Section 1.3. Let (T, E) be a Toeplitz matricial probability space over B, induced by (A, φ). In this section, we will compute operator-valued moments and operator-valued cumulants of a single B-valued random variable

$$A = (a_1, ..., a_N) \equiv \begin{pmatrix} a_1 & a_2 & \cdots & a_N \\ & a_1 & \ddots & \vdots \\ & & \ddots & a_2 \\ O & & & a_1 \end{pmatrix} \in (T, E).$$

We will compute the moment series of A, $M_A(z)$ and the R-transform of A, $R_A(z)$, in Θ_B^1. To do that, we have to compute the n-th moment of A and n-th cumulant of A, as n-th coefficient of M_A and R_A, respectively, for $n \in \mathbb{N}$. Recall that, since $C_T(B) = B$, we can define M_A and R_A by trivial moment series of A

and trivial R-transform of A, in the sense of Section 1.2 (See Section 2.2 and 2.3). i.e

$$M_A(z) = \sum_{n=1}^{\infty} E\left(A^n\right)\, z^n$$

and

$$R_A(z) = \sum_{n=1}^{\infty} K_n\left(A, ..., A\right)\ z^n,$$

where $E\left((a_1, ..., a_N)\right) = \left(\varphi(a_1), ..., \varphi(a_n)\right)$, for all $(a_1, ..., a_N) \in (T, E)$ and K_n is induced by E, in the sense of Section 2.3.

Consider the general expression of A^n, for $n \in \mathbb{N}$. Similar to the previous section, we can compute that ;

$$A^2 = (a_1, ..., a_N) \cdot (a_1, ..., a_N)$$

$$\equiv \begin{pmatrix} a_1 & a_2 & \cdots & a_N \\ & a_1 & \ddots & \vdots \\ & & \ddots & a_2 \\ O & & & a_1 \end{pmatrix} \begin{pmatrix} a_1 & a_2 & \cdots & a_N \\ & a_1 & \ddots & \vdots \\ & & \ddots & a_2 \\ O & & & a_1 \end{pmatrix}$$

$$= \begin{pmatrix} a_1^2 & \sum_{k=1}^{2} a_k a_{(2+1)-k} & \cdots & \sum_{k=1}^{N} a_k a_{(N+1)-k} \\ & a_1^2 & \ddots & \vdots \\ & & \ddots & \sum_{k=1}^{2} a_k a_{(2+1)-k} \\ & & & a_1^2 \end{pmatrix}$$

$$\equiv \left(a_1^2,\ \sum_{k=1}^{2} a_k a_{(2+1)-k},\ ...,\ \sum_{k=1}^{N} a_k a_{(N+1)-k}\right)$$
$$\overset{denote}{=} \left(P_1^{(2)}, P_2^{(2)}, ..., P_N^{(2)}\right).$$

Inductively, we have that

$$P_j^{(1)} = a_j,$$

and

$$P_j^{(n)} = \sum_{k=1}^{j} P_k^{(n-1)} a_{(j+1)-k},$$

for all $j = 1, ..., N$.

Thus we can compute the n-th moment of A ;

$$E\left(A^n\right) = E\left((P_1^{(n)}, ..., P_N^{(n)})\right) = \left(\varphi(P_1^{(n)}), ..., \varphi(P_N^{(n)})\right) \in B.$$

For example, if $N = 2$, then we have that

$P_1^{(1)} = a_1$ and $P_2^{(1)} = a_2$

$P_1^{(2)} = a_1^2$ and $P_2^{(2)} = a_1a_2 + a_2a_1$

$P_1^{(3)} = a_1^3$ and $P_2^{(3)} = a_1^2a_2 + a_1a_2a_1 + a_2a_1^2 = \sum_{k=1}^{2} P_k^{(2)} a_{(2+1)-k}$

etc, and hence

$$\begin{aligned} E\left(A^3\right) &= \left(\varphi(a_1^3), \varphi\left(a_1^2a_2 + a_1a_2a_1 + a_2a_1^2\right)\right) \\ &= \left(\varphi(a_1^3),\ \varphi(a_1^2a_2) + \varphi(a_1a_2a_1) + \varphi(a_2a_1^2)\right) \end{aligned}$$

and
$K_3\left(A, A, A\right) = \left(k_3(a_1, a_1, a_1),\ k_3(a_1, a_1, a_2) + k_3(a_1, a_2, a_1) + k_3(a_2, a_1, a_1)\right).$

Proposition 3.1. *Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace. Let (T, E) be a Toeplitz matricial probability space over B and let $A = (a_1, ..., a_N) \in (T, E)$ be a B-valued random variable. Then*

$$M_A(z) = \sum_{n=1}^{\infty} \left(\varphi(P_1^{(n)}, ..., P_N^{(n)})\right) z^n$$

and

$$R_A(z) = \sum_{n=1}^{\infty} \left(k_n(Q_1^{(n)}), ..., k_n(Q_N^{n})\right) z^n,$$

in Θ_B^1, where

$$Q_j^{(n)} = \sum_{k=1}^{j} \left(Q_k^{(n-1)},\ a_{(j+1)-k}\right) \in \underbrace{A \times .... \times A}_{n-times},$$

with $Q_j^{(1)} = a_j$, for all $j = 1, ..., N$. $\square$

Theorem 3.2. *Let $B = \mathcal{C}^N$ be a Toeplitz matricial algebra and let (A, φ) be a NCPSpace in the sense of Section 1.3. Let (T, E) be a Teoplitz matricial probability space over B, induced by (A, φ) and let $A = (a_1, ..., a_N) \in (T, E)$ be a B-valued random variable, with $a_1, ..., a_N \in (A, \varphi)$ are random variables. If $\{a_1\}, ..., \{a_N\}$ are free in (A, φ) (i.e $\{a_1, ..., a_N\}$ is a free family in (A, φ)), then*

$$R_A(z) = (\varphi(a_1), ..., \varphi(a_N))\, z + b_2 z^2 + \sum_{n=3}^{\infty} \left(k_n(\underbrace{a_1, ..., a_1}_{n-times}), ..., 0 \right) z^n,$$

in Θ_B^1, where $b_2 \in B$ such that

$$b_2 = \begin{cases} \left(k_2(a_1, a_1), 0, k_2(a_2, a_2), 0, ..., 0, k_2(a_{\frac{N}{2}},\ a_{\frac{N}{2}}), 0\right) & N \textit{ is even} \\ \\ \left(k_2(a_1, a_1), 0, k_2(a_2, a_2), 0, ..., 0, k_2(a_{\frac{N-1}{2}},\ a_{\frac{N-1}{2}})\right) & N \textit{ is odd.} \end{cases}$$

Proof. By the previous proposition, we have n-th coefficient of $R_A(z)$,

$$K_n\left(\underbrace{A, ..., A}_{n-times}\right) = \left(k_n(Q_1^{(n)}), ..., k_n(Q_N^{(n)})\right).$$

We need to consider $Q_j^{(n)}$, for $j = 1, ..., N$. Since $Q_j^{(n)}$ is gotten from $P_j^{(n)}$, $j = 1, ..., N$, we will observe $P_j^{(n)}$;

$$P_j^{(1)} = a_j,$$

$$P_j^{(2)} = \textstyle\sum_{k=1}^{j} P_k^{(1)} a_{(j+1)-k}$$

and

$$P_j^{(n)} = \textstyle\sum_{k=1}^{j} P_k^{(n-1)} a_{(j+1)-k},$$

for all $j = 1, ..., N$. Hence

$$Q_j^{(1)} = P_j^{(1)} = a_j,$$

$$Q_j^{(2)} = \textstyle\sum_{k=1}^{j} \left(Q_k^{(1)},\ a_{(j+1)-k}\right)$$

and

$$Q_j^{(n)} = \textstyle\sum_{k=1}^{j} \left(Q_k^{(n-1)}, a_{(j+1)-k}\right),$$

for all $j = 1, ..., N$. Thus

$$K_1(A) = (k_1(a_1), ..., k_1(a_N)) = (\varphi(a_1), ..., \varphi(a_N)),$$

$$K_2(A, A) = (k_2(a_1, a_1),\ k_2\,(a_1, a_2) + k_2(a_2, a_1),$$
$$..., k_2(a_1, a_N) + k_2(a_2, a_{N-1}) + ... + k_2(a_N, a_1))$$

and

$$K_n(A, ..., A)$$
$$= \left(k_n(a_1, ..., a_1), \textstyle\sum_{k=1}^{2} k_2(Q_k^{(n-1)}, a_{3-k}), ..., \sum_{k=1}^{N} k_n(Q_k^{(n-1)}, a_{(N+1)-k})\right).$$

(i) If $n = 1$, then we have that

$$K_1(A) = (k_1(a_1), k_1(a_2), ..., k_1(a_N)) = (\varphi(a_1), ..., \varphi(a_N)).$$

(ii) If $n \geq 3$, then, since $\left(Q_k^{(n-1)}, a_{(j+1)-k}\right)$ is a mixed tuple over $\{a_1, ..., a_N\}$, for all $j = 1, ..., N$ and $k \in \{\boxed{2}, ..., N\}$, we have that

$$K_n\left(\underbrace{A, ..., A}_{n-times}\right) = \left(k_n\left(\underbrace{a_1, ..., a_1}_{n-times}\right), 0, ..., 0\right)$$

$$\equiv \begin{pmatrix} k_n(a_1, ..., a_1) & & & O \\ & k_1(a_1, ..., a_1) & & \\ & & \ddots & \\ O & & & k_1(a_1, ..., a_1) \end{pmatrix} \in B.$$

(iii) If $n = 2$, then we have following two cases ;

<Case 1> Suppose that $N = 2k$, for $k \in \mathbb{N}$. Then

$$K_2(A, A) = (k_2(a_1, a_1),\ k_2\,(a_1, a_2) + k_2(a_2, a_1),$$
$$..., k_2(a_1, a_N) + k_2(a_2, a_{N-1}) + ... + k_2(a_N, a_1))$$

$$= \left(k_2(a_1, a_1), 0, k_2(a_2, a_2), 0, k_2(a_3, a_3), 0, ..., 0, k_2(a_{\frac{N}{2}},\ a_{\frac{N}{2}}), 0\right)$$

$$\equiv \begin{pmatrix} k_2(a_1, a_1) & 0 & & \cdots & k_2(a_{\frac{N-1}{2}},\ a_{\frac{N-1}{2}}) & 0 \\ & k_2(a_1, a_1) & 0 & & * & k_2(a_{\frac{N-1}{2}},\ a_{\frac{N-1}{2}}) \\ & & & \ddots & & \vdots \\ & & \ddots & \ddots & & \\ & & & & & 0 \\ O & & & & & k_2(a_1, a_1) \end{pmatrix}_{N \times N}$$

<Case 2> Suppose that $N = 2k - 1$, for $k \in \mathbb{N}$. Then

$$K_2(A, A) = \left(k_2(a_1, a_1), 0, k_2(a_2, a_2), 0, ..., 0, k_2(a_{\frac{N-1}{2}},\ a_{\frac{N-1}{2}})\right)$$

$$\equiv \begin{pmatrix} k_2(a_1, a_1) & 0 & & \cdots & 0 & k_2(a_{\frac{N-1}{2}},\, a_{\frac{N-1}{2}}) \\ & k_2(a_1, a_1) & & & * & 0 \\ & & & & \ddots & \vdots \\ & & \ddots & \ddots & & \\ & & & & & 0 \\ O & & & & & k_2(a_1, a_1) \end{pmatrix}_{N \times N}$$

By (i), (ii) and (iii), we have that

$$\begin{aligned} R_A(z) = {} & (\varphi(a_1), ..., \varphi(a_N))\, z \\ & + \Big(k_2(a_1, a_1), 0, k_2(a_2, a_2), 0, ..., 0, k_2(a_{\frac{N}{2}},\, a_{\frac{N}{2}}), 0\Big)\, z^2 \\ & + (k_3(a_1, a_1, a_1), 0, ..., 0)\, z^3 + ... \end{aligned}$$

$$= (\varphi(a_1), ..., \varphi(a_N))\, z + \Big(k_2(a_1, a_1), 0, k_2(a_2, a_2), 0, ..., 0, k_2(a_{\frac{N}{2}},\, a_{\frac{N}{2}}), 0\Big)\, z^2 + \sum_{n=3}^{\infty} \left(k_n \left(\underbrace{a_1, ..., a_1}_{n-times} \right), 0, ..., 0 \right) z^n,$$

if N is even, and

$$\begin{aligned} R_A(z) = {} & (\varphi(a_1), ..., \varphi(a_N))\, z \\ & + \Big(k_2(a_1, a_1), 0, k_2(a_2, a_2), 0, ..., 0, k_2(a_{\frac{N-1}{2}},\, a_{\frac{N-1}{2}})\Big)\, z^2 \\ & + (k_3(a_1, a_1, a_1), 0, ..., 0)\, z^3 + ... \end{aligned}$$

$$= (\varphi(a_1), ..., \varphi(a_N))\, z + \Big(k_2(a_1, a_1), 0, k_2(a_2, a_2), 0, ..., 0, k_2(a_{\frac{N-1}{2}},\, a_{\frac{N-1}{2}})\Big)\, z^2 + \sum_{n=3}^{\infty} \left(k_n \left(\underbrace{a_1, ..., a_1}_{n-times} \right), 0, ..., 0 \right) z^n,$$

if N is odd. ∎

3.2. $\mathcal{C}^N$-**Evenness.**

First, we will consider the B-valued evenness, where B is an arbitrary unital algebra. Let B be a unital algebra and let (A, E) be a NCPSpace over B. In [12], we defined B-valued evenness and we observed some properties of it. In this section, we want to characterize the $\mathcal{C}^N$-evenness, where $B = \mathcal{C}^N$, a Toeplitz matricial algebra, for $N \in \mathbb{N}$. Throughout this section, fix $N \in \mathbb{N}$.

Definition 3.1. *Let B be a unital algebra and let (A, E) be a NCPSpace over B. We say a B-valued random variable, $x \in (A, E)$, is an (B-valued) even element if x satisfies the following cumulant relation ;*

$$K_n \left(\underbrace{x, ..., x}_{n-times} \right) = 0_B, \text{ whenever } n \in \mathbb{N} \text{ is odd,}$$

where

$$K_n(x, ..., x) = c^{(n)} \left(x \otimes b_{i_2} x \otimes ... \otimes b_{i_n} x \right),$$

is a n-th cumulant induced by a B-functional, $E : A \to B$, for $b_{i_2}, ..., b_{i_n} \in B$, arbitrary.

In [12], we used the alternating definition for evenness as follows ;

$$x \in (A, E) \text{ is even} \Leftrightarrow \text{moments } E\left(x b_2 x ... b_n x\right) = 0_B, \text{ whenever } n \text{ is odd.}$$

Also, in [12], we observed that our definition with respect to cumulnats and the above definition with respect to moments are equivalent.

Theorem 3.3. *(Also see [12])Let B be a unital algebra and let (A, E) be a NCPSpace over B. If a B-valued random variable, $x \in (A, E)$, is even, then*

$$R_x(z) = \sum_{n=1}^{\infty} K_{2n} \left(\underbrace{x, ..., x}_{2n-times} \right) z^n \in \Theta_B^1$$

with

$$K_{2n} \left(\underbrace{x, ..., x}_{2n-times} \right) = \sum_{\pi \in NC^{(even)}(2n)} \widehat{E}(\pi) \left(x \otimes b_2 x \otimes ... \otimes b_{2n} x \right) \mu(\pi, 1_{2n}),$$

where

$$NC^{(even)}(2n) \overset{def}{=} \{\theta \in NC(2n) : V \in \theta \Leftrightarrow |V| \text{ is even}\}.$$

Proof. Remark that x is even if and only if that all odd moments vanish. Suppose that $\pi \in NC(2n)$ and assume that there exists at least one $V \in \pi$ satisfies that $|V| \in \mathbb{N}$ is an odd number (i.e a block, $V \in \pi$, is an odd block).

(i) Suppose that such odd block V is contained in $\pi(o)$. Then

$$\begin{aligned}
&\widehat{E}(\pi)\,(x \otimes b_2 x \otimes ... \otimes b_n x) \\
&\quad = \prod_{W \in \pi(o)} \widehat{E}(\pi \mid_W)(x \otimes b_2 x \otimes ... \otimes b_n x) \\
&\quad = \left(\widehat{E}(\pi \mid_V)(x \otimes b_2 x \otimes ... \otimes b_n x)\right) \\
&\qquad\qquad \cdot \left(\prod_{W \neq V \in \pi(o)} \widehat{E}(\pi \mid_W)(x \otimes b_2 x \otimes ... \otimes b_n x)\right) \\
&\quad = 0_B.
\end{aligned}$$

(ii) Suppose that such odd block V is contained in $\pi(i)$. Then

$$\begin{aligned}
&\widehat{E}(\pi)\,(x \otimes b_2 x \otimes ... \otimes b_n x) \\
&\quad = \prod_{W \in \pi(o)} \widehat{E}(\pi \mid_W)(x \otimes b_2 x \otimes ... \otimes b_n x).
\end{aligned}$$

Assume that the odd block $V = (v_1, ..., v_k) \in \pi(i)$ is an inner block with its outer block $W_0 = (w_1, ..., w_p) \in \pi(o)$ and assume that there exists $j \in \{1, ..., p\}$ such that $w_j < v_q < w_{j+1}$, for all $q = 1, ..., k$. Then the right hand side of the above equation is

$$\begin{aligned}
&= \left(\widehat{E}(\pi \mid_{W_0})(x \otimes b_2 x \otimes ... \otimes b_n x)\right) \\
&\qquad\qquad \cdot \left(\prod_{W \neq W_0 \in \pi(o)} \widehat{E}(\pi \mid_W)(x \otimes b_2 x \otimes ... \otimes b_n x)\right) \\
&= \left(E^{(p)}(b_{w_1} x \otimes ... \otimes b_j x \otimes E^{(k)}(b_{v_1} x \otimes ... \otimes b_{v_k} x) b_{j+1} x \otimes ... \otimes b_p x)\right) \\
&\qquad\qquad \cdot \left(\prod_{W \neq W_0 \in \pi(o)} \widehat{E}(\pi \mid_W)(x \otimes b_2 x \otimes ... \otimes b_n x)\right) \\
&= 0_B.
\end{aligned}$$

So, by (i) and (ii), if $\pi \in NC(2n)$ contains an odd block, then $\widehat{E}(\pi)(x \otimes b_2 x \otimes ... \otimes b_{2n} x)$ vanishs. So, $K_{2n}(x, ..., x)$ is determind by

$$K_{2n}(x, ..., x) = \sum_{\pi \in NC(2n)} \widehat{E}(\pi)\,(x \otimes b_2 x \otimes ... \otimes b_n x)\,\mu(\pi, 1_{2n})$$

$$= \sum_{\pi \in NC^{(even)}(2n)} \widehat{E}(\pi)\,(x \otimes b_2 x \otimes ... \otimes b_n x)\,\mu(\pi, 1_{2n})$$

$$+\sum_{\pi\in NC(2n)\,\setminus NC^{(even)}(2n)}\widehat{E}(\pi)\,(x\otimes b_2x\otimes ...\otimes b_nx)\,\mu(\pi,1_{2n})$$

$$=\sum_{\pi\in NC^{(even)}(2n)}\widehat{E}(\pi)\,(x\otimes b_2x\otimes ...\otimes b_nx)\,\mu(\pi,1_{2n})+0_B. \ \blacksquare$$

From now, let $B=\mathcal{C}^N$ be a Toeplitz matricial algebra. We will characterize the Toeplitz matricial evenness. Recall that, since $C_T(B)=B$, we just defined Toeplitz matricial moments and Toeplitz matricial cumulants as trivial moments and trivial cumulants.

Theorem 3.4. *Let $B=\mathcal{C}^N$ be a Toeplitz matricial algebra and let (A,φ) be a NCPSpace in the sense of Section 1.3. Let (T,E) be a Toeplitz matricial probability space over B, induced by (A,φ). A B-valued random variable $X=(x_1,...,x_N)\in (T,E)$ is (B-valued) even if and only if $E\left(\underbrace{X,.....,X}_{n-times}\right)=0_B$, whenever n is odd. Equivalently, $X=(x_1,...,x_N)\in(T,E)$ is even if and only if*

$$\varphi(x_1)=...=\varphi(x_N)=0,\ in\ \mathbb{C}$$

and

$$\textstyle\sum_{k=1}^{j}\varphi\left(P_k^{(2l)}x_{(j+1)-k}\right)=0,\ in\ \mathbb{C},$$

for all $j=1,...,N$, where

$$X^{2l+1}=(x_1,...,x_N)^{2l+1}=\left(P_1^{(2l+1)},...,P_N^{(2l+1)}\right),\ \forall\ l\in\mathbb{N}.$$

Proof. By an alternating definition of amalgamated evenness, we have that $X=(x_1,...,x_N)\in(T,E)$ is even if and only if all odd moments vanishs. So, $E\left(X^n\right)=0_B$, whenever $n\in\mathbb{N}$ is odd. Suppose that n is odd. Then

$$\begin{aligned}E\left(X^n\right)&=E\left((P_1^{(n)},....,P_N^{(n)})\right)\\&=\left(\varphi(P_1^{(n)}),....,\varphi(P_N^{(n)})\right)\\&=0_B=(0,...,0),\end{aligned}$$

with

$$P_j^{(n)}=\textstyle\sum_{k=1}^{j}P_k^{(n-1)}x_{(j+1)-k},$$

where

$$P_j^{(1)}=x_j\quad\text{and}\quad P_j^{(2)}=\textstyle\sum_{k=1}^{j}x_kx_{(j+1)-k},$$

in (A,φ), for all $j=1,...,N$. Hence we have that

$$\varphi\left(P_j^{(n)}\right) = \varphi\left(\sum_{k=1}^{j} P_j^{(n-1)} x_{(j+1)-k}\right) = 0 \in \mathbb{C},$$

for all $j = 1, ..., N$.

(i) Let $n = 1$. Then

$$\varphi\left(P_j^{(1)}\right) = 0 \in \mathbb{C}, \text{ for } j = 1, ..., N,$$

equivalently,

$$\varphi(x_j) = 0 \in \mathbb{C}, \text{ for all } j = 1, ..., N.$$

(ii) Suppose that $n = 2l + 1$, for $l \in \mathbb{N}$. Then

$$\begin{aligned}\varphi\left(P_j^{(2l+1)}\right) &= \varphi\left(\sum_{k=1}^{j} P_j^{(2l)} x_{(j+1)-k}\right)\\ &= \sum_{k=1}^{j} \varphi\left(P_j^{(2l)} x_{(j+1)-k}\right) = 0 \quad \in \mathbb{C},\end{aligned}$$

for all $j = 1, ..., N$. ∎

Example 3.1. *Let $N = 2$. Suppose that a $B = \mathcal{C}^2$-valued random variable, $X = (x_1, x_2) \in (T, E)$, is even, where (T, E) is a Toeplitz matricial probability space over B, induced by a NCPSpace (A, φ). Then we have the following relation ;*

$$\varphi(x_1) = 0 = \varphi(x_2)$$

and

$$\varphi\left(P_1^{(2l+1)}\right) = \varphi\left(x_1^{2l+1}\right) = 0 \quad \text{and} \quad \sum_{k=1}^{2} \varphi\left(P_k^{(2l)} x_{(2+1)-k}\right) = 0,$$

in $\mathbb{C}$, for all $l \in \mathbb{N}$. Note that

$$\begin{aligned}&\sum_{k=1}^{2} \varphi\left(P_k^{(2l)} x_{(2+1)-k}\right) = 0\\ &\qquad\Longleftrightarrow \varphi\left(P_1^{(2l)} x_2\right) + \varphi\left(P_2^{(2l)} x_1\right) = 0\\ &\qquad\Longleftrightarrow \varphi\left(x_1^{2l} \cdot x_2\right) + \varphi\left(P_2^{(2l)} x_1\right) = 0\\ &\qquad\Longleftrightarrow \varphi\left(P_2^{(2l)} x_1\right) = -\varphi\left(x_1^{2l} \cdot x_2\right).\end{aligned}$$

Observe that

$$X = (x_1, x_2) \equiv \begin{pmatrix} x_1 & x_2 \\ 0 & x_1 \end{pmatrix} \in (T, E)$$

and

$$X^{2l+1} = (x_1, x_2)^{2l+1} = \left(P_1^{(2l+1)},\, P_2^{(2l+1)}\right)$$

$$\equiv \begin{pmatrix} P_1^{(2l+1)} & P_2^{(2l+1)} \\ 0 & P_1^{(2l+1)} \end{pmatrix} = \begin{pmatrix} x_1^{2l+1} & P_2^{(2l+1)} \\ 0 & x_1^{2l+1} \end{pmatrix} = \begin{pmatrix} x_1^{2l+1} & P_2^{(2l+1)} \\ 0 & x_1^{2l+1} \end{pmatrix}$$

$$= \begin{pmatrix} x_1^{2l+1} & x_1^{2l} \cdot x_2 + P_2^{(2l)} x_1 \\ 0 & x_1^{2l+1} \end{pmatrix} = \left(x_1^{2l+1},\, x_1^{2l} \cdot x_2 + P_2^{(2l)} \cdot x_1\right) \in (T, E).$$

Therefore, we can get that ;

(i) if $a \in (A, \varphi)$ *is an (scalar-valued) even random variable (in the sense that* $a \in (A, \varphi)$ *is even if and only if all odd scalar-valued moments vanish), then* $(a, 0) \equiv \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \in (T, E)$ *is again an even* $B = \mathcal{C}^2$*-valued random variable.*

(ii) if $a \in (A, \varphi)$ *is an (scalar-valued) even random variable, then* $(a, a) \equiv \begin{pmatrix} a & a \\ 0 & a \end{pmatrix} \in (T, E)$ *is again an even* $B = \mathcal{C}^2$*-valued random variable.*

(iii) if $a_1, a_2 \in (A, \varphi)$ *are* ***free*** *(scalar-valued) even random variables and if they are identically distributed, then* $(a_1, a_2) \equiv \begin{pmatrix} a_1 & a_2 \\ 0 & a_1 \end{pmatrix} \in (T, E)$ *is again an even* $B = \mathcal{C}^2$*-valued random variable, by (ii).*

Corollary 3.5. *Let* $B = \mathcal{C}^N$ *be a Toeplitz matricial algebra and let* (A, φ) *be a NCPSpace in the sense of Section 1.3. Let* (T, E) *be a Toeplitz matricial probability space over* B.

(i) If $x \in (A, \varphi)$ *is an even (scalar-valued) random variable (in the sense of Nica, See [2]), then* $X = (x, 0, ..., 0) \in (T, E)$ *is an even* B*-valued random variable.*

(ii) If $x \in (A, \varphi)$ *is an even (scalar-valued) random variable, then* $X = \left(\underbrace{x, ..., x}_{N-times}\right) \in (T, E)$ *is an even* B*-valued random variable.*

(iii) If $x_1, ..., x_N \in (A, \varphi)$ *are* ***free*** *even (scalar-valued) random variables and if they are identically distributed, then* $X = (x_1, ..., x_N) \in (T, E)$ *is an even* B*-valued random variable.*

Proof. (i) We have that

$$P_1^{(n)} = x^n \text{ and } P_j^{(n)} = 0_B \in (A, \varphi), \text{ for all } j = 2, ..., N.$$

Since $x \in (A, \varphi)$ is even, all odd scalar-valued moments vanish. So,

$$E(X^n) = E\left(x^n, 0_B, ..., 0_B\right)$$

$$= \begin{cases} (\varphi(x^n), 0, ..., 0) \in B & \text{if } n \text{ is even} \\ \\ (0, ..., 0) = 0_B & \text{if } n \text{ is odd} \end{cases}$$

$$= \begin{cases} E(X^n) & \text{if } n \text{ is even} \\ \\ 0_B & \text{if } n \text{ is odd.} \end{cases}$$

Thus X is even, in (T, E).

(ii) It is easy to see that

$$X^n = (x, ..., x)^n$$

$$\equiv \begin{pmatrix} x & x & \cdots & x \\ & x & \ddots & \vdots \\ & & \ddots & x \\ O & & & x \end{pmatrix}^n = \begin{pmatrix} x^n & \alpha_2 x^n & \cdots & \alpha_N\, x^n \\ & x^n & \ddots & \vdots \\ & & \ddots & \alpha_2\, x^n \\ O & & & x^n \end{pmatrix}$$

$$\equiv (x^n, \alpha_2\, x^n, \alpha_3\, x^n, ..., \alpha_N\, x^n) \in (T, E),$$

where $\alpha_2, ..., \alpha_N \in \mathbb{C}$ are arbitrary for each $n \in \mathbb{N}$. So, if n is odd, then

$$E\left(X^n\right) = E\left(x^n, \alpha_2\, x^n, \alpha_3\, x^n, ..., \alpha_N\, x^n\right)$$

$$= (\varphi(x^n),\ \alpha_2\ \cdot \varphi(x^n), ..., \alpha_N\ \cdot \varphi(x^n))$$

$$= \begin{cases} (\varphi(x^n),\ \alpha_2\ \cdot \varphi(x^n), ..., \alpha_N\ \cdot \varphi(x^n)) & \text{if } n \text{ is even} \\ \\ (0, ..., 0) = 0_B & \text{if } n \text{ is odd,} \end{cases}$$

by the (scalar-valued) evenness of $x \in (A, \varphi)$. Therefore, $X = \left(\underbrace{x, ..., x}_{N-times}\right) \in (T, E)$ is even.

(iii) Since $x_1, ..., x_N$ are free each other, by applying the Möbius inversion, we can get the result, similar to (ii). ∎

3.3. Toeplitz Matricial Probability Spaces Induced by A Compressed NCPSpace.

In [3], Speicher considered a (scalar-valued) compressed NCPSpace and compressed cumulants. And, in [2], Nica observed the compressed (scalar-valued) R-transforms. In [32], we observed an opreator-valued version of them. In this section, we will consider the relation between Toeplitz matricial probability space (T_N, E_N), induced by a NCPSpace (A, φ) and Toeplitz matricial probability space $(T_N^p,\, E_N^p)$, induced by a compressed NCPSpace $\left(pAp, \frac{1}{\varphi(p)}\varphi \mid_{pAp}\right)$ of (A, φ), where $p \in A$ is a projection satisfying $\varphi(p) \neq 0$. By Speicher and Nica, a compressed NCPSpace $\left(pAp, \varphi_p\right)$, where $\varphi_p = \frac{1}{\varphi(p)}\varphi \mid_{pAp}$, is again a NCPSpace, in the sense of Section 1.3. From now, we will denote a compressed R-transform of random variable $x \in (A, \varphi)$ and compressed cumulants of x, as coefficients of the R-transform by $R_x^{(pAp)}$ and $k_n^{(pAp)}(x, ..., x)$, respectively. It is known the relation between cumulants and compressed cumulants, by Speicher and Nica. Also, in [12], if B is a unital algebra and (A, E) is a NCPSpace over B, we realized that there is similar relation between compressed operator-valued cumulants and operator-valued cumulants, when we take a projection, p, such that $E(p) \in C_A(B) \cap B_{inv}$, compared with the scalar-valued case.

Definition 3.2. *Let (A, φ) be a NCPSPace in the sense of Section 1.3 and let $p \in A$ be a projection (i.e idempotent, $p^2 = p$, in (A, φ), in short, $p \in A_{pro}$) such that $\varphi(p) \neq 0$. Then $\left(pAp, \varphi_p\right)$, with $\varphi_p = \frac{1}{\varphi(p)}\varphi \mid_{pAp}$, is called a compressed NCPSpace (by $p \in A_{pro}$). As usual, we will denote a Toeplitz matricial probability space, induced by (A, φ), by $(T_N, E_N) \equiv (T, E)$, for $N \in \mathbb{N}$. Denote a Toeplitz matricial probability space induced by the compressed NCPSpace, $\left(pAp, \varphi_p\right)$, by $(T_N^p,\, E_N^p) \equiv (T^p, E^p)$.*

The follwong proposition is proved in [2] and [3] ;

Proposition 3.6. *(See [2] and [3]) Let (A, φ) be a NCPSpace and let $p \in A_{pro}$ satisfy $\varphi(p) \neq 0$, in $\mathbb{C}$. If $x_1, ..., x_s \in (A, \varphi)$ are (scalar-valued) random variables ($s \in \mathbb{N}$), then*

$$R_{x_1,...,x_s}^{(pAp)}(z_1, ..., z_s) = \sum_{n=1}^{\infty} \sum_{i_1,...,i_n \in \{1,...,s\}} \alpha_{i_1,...,i_n}\, z_{i_1} ... z_{i_n},$$

in $\Theta_{\mathbb{C}}^{s}$, *with*

$$\begin{aligned}\alpha_{i_1,...,i_n} &= k_n^{(pAp)}(x_{i_1},...,x_{i_n}) \\ &= \alpha_0^{-1}k_n\left(\alpha_0 x_{i_1},...,\alpha_0 x_{i_n}\right) \\ &= \alpha_0^{n-1}k_n(x_{i_1},...,x_{i_n}),\end{aligned}$$

for all $(i_1,...,i_n) \in \{1,...,s\}^n$, $n \in \mathbb{N}$, *where* $\alpha_0 = \varphi(p) \in \mathbb{C}$. *Here,* k_n *is an* n*-th cumulant and* $k_n^{(pAp)}$ *is a compressed* n*-th cumulant in the sense of Speicher and Nica (See [2] and [3]).* □

Remark 3.1. *More generally, in [32], we showed that if* B *is an arbitrary unital algebra and* (A, E) *is a NCPSpace over* B *and if* $p \in A_{pro}$ *satisfies that*

$$E(p) \in C_A(B) \cap B_{inv},$$

then (pAp, E_p) *is a NCPSpace over* B*, again, where* $E_p = (E(p))^{-1} E \mid_{pAp}$*. (i.e* $B \subset pAp \subset A$*) Now, let* $x_1,...,x_s \in (A, E)$ *be* B*-valued random variables* ($s \in \mathbb{N}$) *and denote* ***trivial*** *cumulant multiplicative bimodule map induced by* E *and those induced by* E_p *by* K_n *and* $K_n^{(pAp)}$*, respectively. Then we have that*

$$\begin{aligned}K_n^{(pAp)}(px_{i_1}p,...,px_{i_n}p) &= c_{E_p}^{(n)}\left(px_{i_1}p \otimes px_{i_2}p \otimes ... \otimes px_{i_n}p\right) \\ &= b_0^{-1}c_E^{(n)}\left(b_0x_{i_1} \otimes b_0x_{i_2} \otimes ... \otimes b_0x_{i_n}\right) \\ &= K_n^{symm(b_0)}\left(x_{i_1},...,x_{i_n}\right),\end{aligned}$$

where $b_0 = E(p) \in B$*. Therefore, we can get that*

$$R_{px_1p,...,px_sp}^{(pAp)\,:\,t}(z_1,...,z_s) = R_{x_1,...,x_s}^{symm(E(p))}(z_1,...,z_s),$$

in Θ_B^s.

A Toeplitz probability space (T_N^p, E_N^p), induced by a compressed NCPSpace (pAp, φ_p), can be understood as an algebra that

$$T_N^p = A\lg\left(\{(px_1p,\ px_2p, ..., px_Np) \in (T_N, E_N) : x_j \in (A,\varphi),\ j = 1,...,N\}\right),$$

where (T_N, E_N) is a Toeplitz matricial probability space induced by (A, φ). i.e if $X = (x_1',...,x_N') \in (T_N^p, E_N^p)$, then there exists $x_1^{(i)},...,x_N^{(i)} \in (A,\varphi)$, $(i \in \mathbb{N})$ such that

$$(x_1',...,x_N') = \textstyle\sum_{i=1}^{\infty}\beta_i \cdot (px_1^{(i)}p,...,px_N^{(i)}p),$$

where $\beta_i \in \mathbb{C}$ and, for each $i \in \mathbb{N}$,

$$\beta_i \left(px_1^{(i)}p, ..., px_N p^{(i)}\right)$$

$$\equiv \begin{pmatrix} \beta_i & & & O \\ & \beta_i & & \\ & & \ddots & \\ O & & & \beta_i \end{pmatrix} \begin{pmatrix} px_1^{(i)}p & px_2^{(i)}p & \cdots & px_N^{(i)}p \\ & px_1^{(i)}p & \ddots & \vdots \\ & & \ddots & px_2^{(i)}p \\ O & & & px_1^{(i)}p \end{pmatrix}$$

$$= \left(\beta_i \cdot px_1^{(i)}p, ..., \beta_i \cdot px_N p^{(i)}\right).$$

So, by the above setting, we have that ;

Theorem 3.7. *Let (T_N^p, E_N^p) be a Toeplitz matricial probability space, induced by a compressed NCPSpace, $\left(pAp, \varphi_p\right)$, of a NCPSpace (A, φ), with $p \in A_{pro}$ such that $\varphi(p) \neq 0$. Let $X_1 = (x_1^{(1)}, ..., x_N^{(1)}), ..., X_s = (x_1^{(s)}, ..., x_N^{(s)}) \in (T_N, E_N)$ be $\mathcal{C}^N$-valued random variables ($s \in \mathbb{N}$) and let $X_1' = (px_1^{(1)}p, ..., px_N^{(1)}p), ..., X_s' = (px_1^{(s)}p, ..., px_N^{(s)}p) \in (T_N^p, E_N^p)$ be $\mathcal{C}^N$-valued random variables. Then*

$$R_{X_1', ..., X_s'}^{(T_N^p)}(z_1, ..., z_s)$$

$$= \sum_{n=1}^{\infty} \sum_{i_1, ..., i_n \in \{1, ..., s\}} \alpha_0^{n-1} K_n(X_{i_1}, ..., X_{i_n})\, z_{i_1} ... z_{i_n},$$

where $\alpha_0 = \varphi(p) \in C$ and K_n's are cumulant multiplicative bimodule map induced by $E_N : T_N \to \mathcal{C}^N$.

Proof. Consider $(i_1, ..., i_n)$-th coefficient of $R_{X_1', ..., X_s'}^{(T_N^p)}$;

$$coef_{i_1, ..., i_n}\left(R_{X_1', ..., X_s'}^{(T_N^p)}\right) = K_n^{(T_N^p)}\left(X_{i_1}', ..., X_{i_n}'\right)$$

$$= K_n^{(T_N^p)}\left((px_1^{(i_1)}p, ..., px_N^{(i_1)}p), ..., (px_1^{(i_n)}p, ..., px_N^{(i_n)}p)\right)$$

$$= \left(k_n^{(pAp)}\left(Q_{1\ :\ p}^{(i_1, ..., i_n)}\right), ..., k_n^{(pAp)}\left(Q_{N\ :\ p}^{(i_1, ..., i_n)}\right)\right)$$

where $Q_{j\ :\ p}^{(i_1, ..., i_n)} \in A^n$ is determined inductively as in Section 2.2 ($j = 1, ..., N$), with respect to $(T_N^p,\ E_N^p)$

$$= \left(\alpha_0^{n-1} k_n(Q_1^{(i_1, ..., i_n)}), ..., \alpha_0^{n-1} k_n(Q_N^{(i_1, ..., i_n)})\right)$$

where $\alpha_0 = \varphi(p) \in \mathbb{C}$ and k_n is a n-th cumulant induced by φ, by the previous proposition. Here, $Q_j^{(i_1,...,i_n)} \in A^n$ is determined inductively as in Section 2.2 $(j = 1, ..., N)$, with respect to $(T_N,\ E_N)$,

$$= \alpha_0^{n-1} \left(k_n(Q_1^{(i_1,...,i_n)}), ..., k_n(Q_N^{(i_1,...,i_n)}) \right)$$

by the previous paragraph

$$= \alpha_n^{n-1} K_n \left(X_{i_1}, ..., X_{i_n} \right). \ \blacksquare$$

By using notations in [12] and [32], we can get that ;

Corollary 3.8. *Under the same hypothesis with the previous theorem,*

$$R_{X_1',...,X_s'}^{(T_N^p)}(z_1, ..., z_s) = R_{X_1,...,X_s}^{symm(b_0)}(z_1, ..., z_s),$$

where

$$b_0 = (\varphi(p),\ 0, ..., 0) \in \mathcal{C}^N$$

$\square$

Remark 3.2. *Let's take a projection* $P = (p, 0_A, ..., 0_A) \in (T_N,\ E_N)$, *where* $p \in (A, \varphi)$ *is a projection (i.e* $p^2 = p$, *in* A*). i.e*

$$P = \begin{pmatrix} p & & & O \\ & p & & \\ & & \ddots & \\ O & & & p \end{pmatrix} \in (T_N, E_N),$$

such that $P^2 = P$, *in* T_N. *Suppose that* $\varphi(p) \neq 0$. *Then*

$$E(P) = E\left(p, 0_A, ..., 0_A\right) = (\varphi(p), 0, ..., 0) \in \mathcal{C}^N.$$

Since $\varphi(p) \neq 0$, $E(P) \in \mathcal{C}^N$ *is invertible. So,* $E(P)$ *satisfies the conditions in Remark 3.1. i.e*

$$E(P) \in C_{T_N}(\mathcal{C}^N)\ \cap\ (\mathcal{C}^N)_{inv}.$$

And, by the previous corollary, we can see that we have the same result with Remark 3.1, when we consider the Toeplitz matricial probability space $(T_N^p,\ E_N^p)$, *induced by* (pAp, φ_p). *i.e this Toeplitz matricial probability space* $(T_N^p,\ E_N^p)$ *is nothing but a compressed (amalgamated) NCPSpace over* $\mathcal{C}^N$, $\left(PT_NP,\ E_N(P)^{-1}E_N\mid_{PT_NP}\right)$.

Corollary 3.9. *Let* (A, φ) *be a NCPSpace, in the sense of Section 1.3 and let* $\left(pAp, \varphi_p\right)$ *be a compressed NCPSpace, by* $p \in A_{pro}$, *with* $\varphi(p) \neq 0$, *in* $\mathbb{C}$. *Let* $(T_N,\ E_N)$ *and* $(T_N^p,\ E_N^p)$ *be Toeplitz matricial probability spaces induced by* (A, φ) *and* $\left(pAp, \varphi_p\right)$, *respectively. Then* $(T_N^p,\ E_N^p)$ *is an amalgamated compressed NCPSpace over* $\mathcal{C}^N$, *by* $P = (p, 0_A, ..., 0_A) \in (T_N,\ E_N)$. *i.e*

$$(T_N^p,\ E_N^p) = \left(PT_NP,\ E_N(P)^{-1}E_N\mid_{PT_NP}\right).$$

□

A Toeplitz matricial probability theory can be a good example of amalgamated free probability theory (See [1] and [12]). Futhermore, since $C_{T_N}(\mathcal{C}^N) = \mathcal{C}^N$, it is easy to deal with. i.e we can avoid a difficulty caused by Insertion property in Amalgamated free probability and hence it is enough to consider trivial R-transforms, in the sense of Section 1.2, (defined as Toeplitz matricial R-transforms, in this paper) when we work for R-transform theory and R-transform calculus, on Toeplitz matricial probability spaces.

References

[1] R. Speicher, Combinatorial Theory of the Free Product with Amalgamation and Operator-Valued Free Probability Theory, AMS Mem, Vol 132 , Num 627 , (1998).

[2] A. Nica, R-transform in Free Probability, IHP course note, available at `www.math.uwaterloo.ca/~anica`.

[3] R. Speicher, Combinatorics of Free Probability Theory IHP course note, available at `www.mast.queensu.ca/~speicher`.

[4] A. Nica, D. Shlyakhtenko and R. Speicher, R-cyclic Families of Matrices in Free Probability, J. of Funct Anal, 188 (2002), 227-271.

[5] A. Nica and R. Speicher, R-diagonal Pair-A Common Approach to Haar Unitaries and Circular Elements, (1995), `www.mast.queensu.ca/~speicher`.

[6] D. Shlyakhtenko, Some Applications of Freeness with Amalgamation, J. Reine Angew. Math, 500 (1998), 191-212.

[7] A. Nica, D. Shlyakhtenko and R. Speicher, R-diagonal Elements and Freeness with Amalgamation, Canad. J. Math. Vol 53, Num 2, (2001) 355-381.

[8] A. Nica, R-transforms of Free Joint Distributions and Non-crossing Partitions, J. of Func. Anal, 135 (1996), 271-296.

[9] D.Voiculescu, K. Dykemma and A. Nica, Free Random Variables, CRM Monograph Series Vol 1 (1992).

[10] D. Voiculescu, Operations on Certain Non-commuting Operator-Valued Random Variables, Astérisque, 232 (1995), 243-275.

[11] D. Shlyakhtenko, A-Valued Semicircular Systems, J. of Funct Anal, 166 (1999), 1-47.

[12] I. Cho, Operator-Valued Free Probability and its Application (preprint).

[13] R. Speicher, A Conceptual Proof of a Basic Result in the Combinatorial Approach to Freeness, `www.mast.queensu.ca/~speicher`.

[14] M. Bożejko, M. Leinert and R. Speicher, Convolution and Limit Theorems for Conditionally Free Random Variables, `www.mast.queensu.ca/~speicher`.

[15] A. Nica, R-diagonal Pairs Arising as Free Off-diagonal Compressions, available at
`www.math.uwaterloo.ca/~anica`.

[16] B.Krawczyk and R.Speicher, Combinatorics of Free Cumulants, available at
`www.mast.queensu.ca/~speicher`.

[17] A. Nica and R.Speicher, A "Fouries Transform" for Multiplicative Functions on Noncrossing Patitions, J. of Algebraic Combinatorics, 6, (1997) 141-160.

[18] P.Śniady and R.Speicher, Continous Family of Invariant Subspaces for R-diagonal Operators, Invent Math, 146, (2001) 329-363.

[19] I.Cho, Amalgamated S-transforms, preprint.

[20] I.Cho, B-Poisson distribution and Amalgamated Conditional Freeness, preprint.

[21] Y. Watatani, Index for C*-subalgebras, Mem of AMS, Vol 83, 424 (1990).

[22] V. Jones, Index for Subfactors, Invent. Math, 72 (1983) 1-25.

[23] V. Jones, Braid Groups, Hecke Algebras and Type II_1-factors, Geometric Methods in Operator Algebra, Pitman Res. Notes 123 (1986) 242-273.

[24] V. Jones, Hecke Algebra Representations of Braid Groups and Link Polynomials, Ann Math, 126 (1987) 335-388.

[25] H. Kosaki, Extension of Jones' Theory on Index to Arbitrary Factors, J. of F.A, 66 (1986), 123-140.

[26] M. Pimsner and S. Popa, Iterating the Basic Construction, Trans. AMS, 310, No 1 (1988), 127-133.

[26] S. Popa, Markov Traces on Universal Jones Algebras and Subfactors of Finite Index. Invent. Math, 111 (1993), 375-405.

[27] I. Cho, Amalgamated R-transform Theory of Index-Finite type Conditional Expectations, preprint.

[28] I. Cho, B-free family of Cuntz Isometries, preprint.

[29] I. Cho, Amalagmated Free Probability Theory and its Application, preprint.

[30] A. Nica, D. Shlyakhtenko, R. Speicher, Operator-Valued Distributions I. Characterizations of Freeness, preprint.

[31] A. Nica, D. Shlyakhtenko, F. Goodman, Free Probability of Type B, preprint.

[32] I. Cho, Compressed Amalgamated R-transform Theory, preprint.

[33] I. Cho, Perturbed R-transform Theory, preprint.

Dep. of Math, Univ. of Iowa, Iowa City, IA, U.S.A
E-mail address: `icho@math.uiowa.edu`